

Package ‘SwissAir’

January 2, 2012

Version 1.1.01

Date 2011-11-09

Title Air Quality Data of Switzerland for one year in 30 min Resolution

Maintainer Rene Locher <rene.locher@zhaw.ch>

Author Rene Locher

Description Ozone, NOx (= Sum of Nitrogenmonoxide and Nitrogendioxide), Nitrogenmonoxide, ambient temperature, dew point, wind speed and wind direction at 3 sites around lake of Lucerne in Central Switzerland in 30 min time resolution for year 2004.

LazyData yes

Depends R(>= 2.13.1)

Suggests IDPmisc(>= 1.1.16)

License GPL (>= 3)

Repository CRAN

Date/Publication 2011-11-09 15:08:37

R topics documented:

AirQual	2
Index	6

AirQual

Air Quality Data of Switzerland

Description

Dataset contains Ozone (=O3), Sum of Nitrogenmonoxide and Nitrogendioxide (=NOx), Nitrogenmonoxid (=NO), ambient temperature (T), dew point (Td), wind speed (WS) and wind direction (WD) at 3 sites around lake of Lucerne in Central Switzerland in 30 min time resolution for the year 2004.

Usage

```
data(AirQual)
```

Format

A data frame with 17568 observations on the following 22 variables.

start start time of observation (GMT+1h, no day saving time)

ad.O3 Mean concentration [ppb] of O3 in ad

ad.NOx Mean concentration [ppb] of NOx in ad

ad.NO Mean concentration [ppb] of NO in ad

ad.WS Wind speed [m/s] in ad

ad.WD Wind direction in ad

ad.T Mean ambient temperature [deg C] in ad

ad.Td Mean dew point [deg C] in ad

lu.O3 Mean concentration [ppb] of O3 in sz

lu.NOx Mean concentration [ppb] of NOx in sz

lu.NO Mean concentration [ppb] of NO in sz

lu.WS Wind speed [m/s] in sz

lu.WD Wind direction in sz

lu.T Mean ambient temperature [deg C] in sz

lu.Td Mean dew point [deg C] in sz

sz.O3 Mean concentration [ppb] of O3 in sz

sz.NOx Mean concentration [ppb] of NOx in sz

sz.NO Mean concentration [ppb] of NO in sz

sz.WS Wind speed [m/s] in sz

sz.WD Wind direction in sz

sz.T Mean ambient temperature [deg C] in sz

sz.Td Mean dew point [deg C] in sz

Details

The 3 sites are

ad Site in Altdorf is located 100 m east of motorway A2 from Bale to Chiasso, on an open field at the beginning of a more than 2000 m deep valley, at 438 m altitude.

lu Site is located in Sedel next to town of Lucerne 35m above and 250m south of motorway A14 on a small hill with free 360 degree panorama at 484 m altitude.

sz Site is located in Schwyz in an aerea of medium density of buildings next to a shopping center.

NO and O3 react in the atmosphere within seconds to NO2. The production and destruction of the sum of O3 and NO2 (=Ox) takes place on a much lower time scale, so that the spatial (cf. `ipairs` output) and temporal (cf. `ilagplot`) correlation of Ox is much more pronounced than the correlation of O3 or NO.

Note

Type of variable start has changed in Version 1.08 from factor to character to save memory.

Source

The data are collected by inNet corporation <http://www.innetag.ch/> on behalf of in-Luft, an association for air quality controle, of the Swiss Cantons Aargau, Luzern, Nidwalden, Obwalden, Schwyz, Uri und Zug <http://www.in-luft.ch/>.

References

Rene Locher, Andreas Ruckstuhl; Plausibilisierung von Ozon, Stickoxiden und PM10: Statistische Methoden zur Effizienz- und Qualitaetssteigerung der Messdatenplausibilisierung; inLuft 2003

See Also

For viewing large datasets see `library(IDPmisc)`

Examples

```
data(AirQual)
str(AirQual)
sapply(AirQual,function(x) sum(is.na(x)))
cbind(min=sapply(AirQual[,-1], min, na.rm=TRUE),
      median=sapply(AirQual[,-1], median, na.rm=TRUE),
      max=sapply(AirQual[,-1], max, na.rm=TRUE))

if(require(IDPmisc)) {

  ## low correlation, density on logarithmic scale
  ipairs(AirQual[,c("ad.O3","lu.O3","sz.O3")],
        ztrans=function(x){x[x<1] <- 1; log2(x)*10})
  ipairs(AirQual[,c("ad.NOx","lu.NOx","sz.NOx")],
        ztrans=function(x){x[x<1] <- 1; log2(x)*10})
  ilagplot(AirQual[,c("ad.O3")],set.lags = 1:9,
```

```

ztrans=function(x){x[x<1] <- 1; log2(x)*10}

Ox <- AirQual[,c("ad.O3","lu.O3","sz.O3")] +
  AirQual[,c("ad.NOx","lu.NOx","sz.NOx")] -
  AirQual[,c("ad.NO","lu.NO","sz.NO")]
names(Ox) <- c("ad","lu","sz")

## high correlation, density on logarithmic scale
ipairs(Ox,
  ztrans=function(x){x[x<1] <- 1; log2(x)*10})
ilagplot(Ox$ad,set.lags = 1:9,
  ztrans=function(x){x[x<1] <- 1; log2(x)*10})

dat <-
  data.frame(month =as.numeric(substr(AirQual$start,4,5)),
    hour = as.numeric(substr(AirQual$start,12,13)),
    WD = AirQual$ad.WD,
    NOx = AirQual$ad.NOx,
    O3 = AirQual$ad.O3,
    Ox = AirQual$ad.O3+AirQual$ad.NOx-AirQual$ad.NO)

med.dayrose <- rose(dat[,c("NOx","O3","Ox")],
  subset= dat$month>4 & dat$month<10,
  cyclVar=dat$hour, n.cyclVar=24, circle=24,
  FUN=median, na.rm=TRUE)

## NOx (= NO+NO2) and ozone (O3) have a distinct diurnal variation of
## concentration, whereas Ox (= NO2+O3) varies only very slightly
grid.newpage()
plot(med.dayrose,
  general = general.control(lwd=2),
  grid =
  grid.control(ray.n = 12,
    circ.n =2,
    circ.sub.n = 2,
    cyclVar.lab = seq(0,by=2,to=22)),
  title = title.control(text=
    "Day Rose of Medians\nduring summer time"))
grid.newpage()
plot(med.dayrose,
  general = general.control(lwd=3),
  grid =
  grid.control(ray.n = 12,
    circ.n =2,
    circ.sub.n = 2,
    cyclVar.lab = seq(0,by=2,to=22)),
  title = title.control(text=
    "Day Rose of Medians\nduring summer time"))

## exploration of upslope (North) downslope (South) wind system
## during summer time in the valley north of Gotthard
ncol <- 4

```

```
if(interactive()) x11(16,12)
grid.newpage()
pushViewport(viewport(layout=grid.layout(nrow=3,ncol=ncol),
                        width=0.98, height=0.98))

for (hour in seq(0,22,2)) {
  windrose <-
    rose(dat$WD, cyclVar = dat$WD, circle = 360, n.cyclVar = 32,
        subset = dat$hour>=hour & dat$hour<hour+2 &
                dat$month>4 & dat$month<10,
        FUN = function(x) sum(!is.na(x)),
        warn = FALSE)

  pushViewport(viewport(layout.pos.col=(hour/2)%%ncol+1,
                        layout.pos.row=(hour/2)%%ncol+1))
  plot(windrose,
       general=general.control(
         lwd=3),
       grid=grid.control(
         circ.r = seq(0,150,50),
         circ.sub.r = seq(25,150,25),
         circ.between = 0,
         circ.cex = 0.6,
         cyclVar.cex = 1,
         ray.lim=c(0,150)),
       title=title.control(
         text=paste(hour,"o'clock"),
         between=0.3, cex = 1.5),
       key=key.control(title="hour of day"))
  print(hour)
  popViewport()
} ## end for

} else print("Package IDPmisc is not available")
```

Index

*Topic **datasets**

AirQual, [2](#)

AirQual, [2](#)

SwissAir (AirQual), [2](#)