

Package ‘hier.part’

February 14, 2012

Version 1.0-3

Date 2008/01/15

Title Hierarchical Partitioning

Author Chris Walsh <cwalsh@unimelb.edu.au> and Ralph Mac Nally
<Ralph.MacNally@sci.monash.edu.au>

Depends gtools

Description Variance partition of a multivariate data set

Maintainer Chris Walsh <cwalsh@unimelb.edu.au>

License GPL

Repository CRAN

Date/Publication 2008-01-15 07:19:31

R topics documented:

all.regs	2
combos	3
hier.part	4
hier.part.data	6
partition	7
rand.hp	9
Index	12

`all.regs`*Goodness of fit measures for a regression hierarchy*

Description

Calculates goodness of fit measures for regressions of a single dependent variable to all combinations of N independent variables

Usage

```
all.regs(y, xcan, family = "gaussian", gof = "RMSPE", print.vars = FALSE)
```

Arguments

<code>y</code>	a vector containing the dependent variables
<code>xcan</code>	a dataframe containing the n independent variables
<code>family</code>	family argument of <code>glm</code>
<code>gof</code>	Goodness-of-fit measure. Currently "RMSPE", Root-mean-square 'prediction' error, "logLik", Log-Likelihood or "Rsqu", R-squared
<code>print.vars</code>	if FALSE, the function returns a vector of goodness-of-fit measures. If TRUE, a data frame is returned with first column listing variable combinations and the second column listing goodness-of-fit measures.

Details

This function calculates goodness of fit measures for the entire hierarchy of models using all combinations of N dependent variables, and returns them as an ordered list ready for input into the function `partition`. This function requires the `gtools` package in the `gregmisc` bundle

Value

<code>gfs</code>	If <code>print.vars</code> is FALSE, a vector of goodness of fit measures for all combinations of independent variables in the hierarchy or, if <code>print.vars</code> is TRUE, a data frame listing all combinations of independent variables in the first column in ascending order, and the corresponding goodness of fit measure for the model using those variables
------------------	---

Author(s)

Chris Walsh <Chris.Walsh@sci.monash.edu.au>.

References

Hatt, B. E., Fletcher, T. D., Walsh, C. J. and Taylor, S. L. 2004 The influence of urban density and drainage infrastructure on the concentrations and loads of pollutants in small streams. *Environmental Management* **34**, 112–124.

Walsh, C. J., Papas, P. J., Crowther, D., Sim, P. T., and Yoo, J. 2004 Stormwater drainage pipes as a threat to a stream-dwelling amphipod of conservation significance, *Austrogammarus australis*, in southeastern Australia. *Biodiversity and Conservation* **13**, 781–793.

See Also

[hier.part](#), [partition](#), [rand.hp](#)

Examples

```
#linear regression of log(electrical conductivity) in streams
#against seven independent variables describing catchment
#characteristics (from Hatt et al. 2004)
data(urbanwq)
env <- urbanwq[,2:8]
all.regs(urbanwq$lec, env, fam = "gaussian", gof = "Rsqu",
print.vars = TRUE)

#logistic regression of an amphipod species occurrence in
#streams against four independent variables describing
#catchment characteristics (from Walsh et al. 2004)
data(amphipod)
env1 <- amphipod[,2:5]
all.regs(amphipod$australis, env1, fam = "binomial",
gof = "logLik", print.vars = TRUE)
```

combos

All combinations of a hierarchy of models of n variables

Description

Lists a matrix of combinations of 1 to n variables in ascending order

Usage

```
combos(n)
```

Arguments

n an integer: the number of variables

Details

Lists hierarchy of all possible combinations of n variables in ascending order, starting with 1 variable, then all combinations of 2 variables, and so on until the one combination with all n variables. This function is used by `all.regs` to structure the models required for hierarchical partitioning. This function requires the `gtools` package in the `gregmisc` bundle.

Value

a list containing

<code>ragged</code>	a matrix with zeroes in empty elements and 1 in column 1, 2 in column 2 ... n in column n for full elements
<code>binary</code>	a matrix as for <code>ragged</code> , but with 1 in all full elements

Author(s)

Chris Walsh <Chris.Walsh@sci.monash.edu.au>

See Also

[all.regs](#)

hier.part

Goodness of fit calculation and hierarchical partitioning

Description

Partitions variance in a multivariate dataset

Usage

```
hier.part(y, xcan, family = "gaussian", gof = "RMSPE", barplot = TRUE)
```

Arguments

<code>y</code>	a vector containing the dependent variables
<code>xcan</code>	a dataframe containing the n independent variables
<code>family</code>	family argument of <code>glm</code>
<code>gof</code>	Goodness-of-fit measure. Currently "RMSPE", Root-mean-square 'prediction' error, "logLik", Log-Likelihood or "Rsqu", R-squared
<code>barplot</code>	If TRUE, a barplot of I and J for each variable is plotted expressed as percentage of total explained variance.

Details

This function calculates goodness of fit measures for the entire hierarchy of models using all combinations of N independent variables using the function `all.regs`. It takes the list of goodness of fit measures and, using the `partition` function, applies the hierarchical partitioning algorithm of Chevan and Sutherland (1991) to return a simple table listing each variable, its independent contribution (I) and its conjoint contribution with all other variables (J).

Note earlier versions of `hier.part` (<1.0) produced a matrix and barplot of percentage distribution of effects as a percentage of the sum of all Is and Js, as portrayed in Hatt et al. (2004) and Walsh et al. (2004). This version plots the percentage distribution of independent effects only. The sum of Is equals the goodness of fit measure of the full model minus the goodness of fit measure of the null model.

The distribution of joint effects shows the relative contribution of each variable to shared variability in the full model. Negative joint effects are possible for variables that act as suppressors of other variables (Chevan and Sutherland 1991).

At this stage, the partition routine will not run for more than 12 independent variables. This function requires the `gtools` package in the `gregmisc` bundle

Value

a list containing

<code>gfs</code>	a data frame or vector listing all combinations of independent variables in the first column in ascending order, and the corresponding goodness of fit measure for the model using those variables
<code>IJ</code>	a data frame of I, the independent and J the joint contribution for each independent variable
<code>I.perc</code>	a data frame of I as a percentage of total explained variance

Note

The function produces a minor rounding error for analyses with more than 9 independent variables. To check if this error affects the inference from an analysis, run the analysis several times with the variables entered in a different order. There are no known problems for analyses with 9 or fewer variables.

Author(s)

Chris Walsh <Chris.Walsh@sci.monash.edu.au> using c and fortran code written by Ralph MacNally <Ralph.MacNally@sci.monash.edu.au>.

References

- Chevan, A. and Sutherland, M. 1991 Hierarchical Partitioning. *The American Statistician* **45**, 90–96.
- Hatt, B. E., Fletcher, T. D., Walsh, C. J. and Taylor, S. L. 2004 The influence of urban density and drainage infrastructure on the concentrations and loads of pollutants in small streams. *Environmental Management* **34**, 112–124.

Mac Nally, R. 2000 Regression and model building in conservation biology, biogeography and ecology: the distinction between and reconciliation of 'predictive' and 'explanatory' models. *Biodiversity and Conservation* **9**, 655–671.

Walsh, C. J., Papas, P. J., Crowther, D., Sim, P. T., and Yoo, J. 2004 Stormwater drainage pipes as a threat to a stream-dwelling amphipod of conservation significance, *Austrogammarus australis*, in south-eastern Australia. *Biodiversity and Conservation* **13**, 781–793.

See Also

[all.regs](#), [partition](#), [rand.hp](#)

Examples

```
#linear regression of log(electrical conductivity) in streams
#against seven independent variables describing catchment
#characteristics (from Hatt et al. 2004)
data(urbanwq)
env <- urbanwq[,2:8]
hier.part(urbanwq$lec, env, fam = "gaussian", gof = "Rsqu")

#logistic regression of an amphipod species occurrence in
#streams against four independent variables describing
#catchment characteristics (from Walsh et al. 2004)
data(amphipod)
env1 <- amphipod[,2:5]
hier.part(amphipod$australis, env1, fam = "binomial", gof = "logLik")
```

hier.part.data

Example data for hier.part

Description

Example data sets for hier.part package.

Usage

```
data(urbanwq)
data(amphipod)
data(chevan)
```

Details

urbanwq.txt

Seven catchment variables (fimp, catchment imperviousness^{0.25}; sconn, drainage connection^{0.5}; sdensep, septic tank density^{0.5}; unsealden, unsealed road density; fcarea, catchment area^{0.25}; selev, elevation^{0.5}; amgeast, longitude) and median baseflow concentrations of three water quality variables in sixteen streams draining 16 independent subcatchments (ldoc, log(dissolved organic carbon); lec, log(electrical conductivity); lnox, log(nitrate/nitrite). Data from Hatt et al. (2004).

amphipod.txt

Presence-absence data for a stream-dwelling amphipod (*australis*) in 58 sites, with four catchment variables (*fimp*, *imperviousness*^{0.25}; *fconn*, *drainage connection*^{0.25}; *densep*, density of septic tanks; *unseal*, unsealed road density). Data from Walsh et al. (2004).

chevan.txt

Chi-squared (*chisq*) and R-squared (*rsqu*) for logistic and linear regression example from Chevan and Sutherland (1991)

Author(s)

Chris Walsh <Chris.Walsh@sci.monash.edu.au>

References

Chevan, A. and Sutherland, M. 1991 Hierarchical Partitioning. *The American Statistician* **45**, 90–96.

Hatt, B. E., Fletcher, T. D., Walsh, C. J. and Taylor, S. L. 2004 The influence of urban density and drainage infrastructure on the concentrations and loads of pollutants in small streams. *Environmental Management* **34**, 112–124.

Walsh, C. J., Papas, P. J., Crowther, D., Sim, P. T., and Yoo, J. 2004 Stormwater drainage pipes as a threat to a stream-dwelling amphipod of conservation significance, *Austrogammarus australis*, in southeastern Australia. *Biodiversity and Conservation* **13**, 781–793.

partition

Hierarchical partitioning from a list of goodness of fit measures

Description

Partitions variance in a multivariate dataset from a list of goodness of fit measures

Usage

```
partition(gfs, pcan, var.names = NULL)
```

Arguments

<i>gfs</i>	an array as outputted by the function <code>all.regs</code> or a vector of goodness of fit measures from a hierarchy of regressions based on <i>pcan</i> variables in ascending order (as produced by function <code>combos</code> , but also including the null model as the first element)
<i>pcan</i>	the number of variables from which the hierarchy was constructed (maximum = 12)
<i>var.names</i>	an array of <i>pcan</i> variable names, if required

Details

This function applies the hierarchical partitioning algorithm of Chevan and Sutherland (1991) to return a simple table listing of each variable, its independent contribution (I) and its conjoint contribution with all other variables (J). The output is identical to the function `hier.part`, which takes the dependent and independent variable data as its input.

Note earlier versions of `partition` (`hier.part < 1.0`) produced a matrix and barplot of percentage distribution of effects as a percentage of the sum of all Is and Js, as portrayed in Hatt et al. (2004) and Walsh et al. (2004). This version plots the percentage distribution of independent effects only. The sum of Is equals the goodness of fit measure of the full model minus the goodness of fit measure of the null model.

The distribution of joint effects shows the relative contribution of each variable to shared variability in the full model. Negative joint effects are possible for variables that act as suppressors of other variables (Chevan and Sutherland 1991).

At this stage, the `partition` routine will not run for more than 12 independent variables. This function requires the `gtools` package in the `gregmisc` bundle

Value

a list containing

<code>gfs</code>	a data frame listing all combinations of independent variables in the first column in ascending order, and the corresponding goodness of fit measure for the model using those variables
<code>IJ</code>	a data frame of I, the independent and J the joint contribution for each independent variable
<code>I.perc</code>	a data frame of I as a percentage of total explained variance
<code>J.perc</code>	a data frame of J as a percentage of sum of all Js

Note

The function produces a minor rounding error for hierarchies constructed from more than 9 variables. To check if this error affects the inference from an analysis, run the analysis several times with the variables entered in a different order. There are no known problems for hierarchies with 9 or fewer variables.

Author(s)

Chris Walsh <Chris.Walsh@sci.monash.edu.au> using c and fortran code written by Ralph MacNally <Ralph.MacNally@sci.monash.edu.au>.

References

- Chevan, A. and Sutherland, M. 1991 Hierarchical Partitioning. *The American Statistician* **45**, 90–96.
- Hatt, B. E., Fletcher, T. D., Walsh, C. J. and Taylor, S. L. 2004 The influence of urban density and drainage infrastructure on the concentrations and loads of pollutants in small streams. *Environmental Management* **34**, 112–124.

See Also

[all.regs](#), [partition](#), [rand.hp](#)

Examples

```
#linear regression of log(electrical conductivity) in streams
#against seven independent variables describing catchment
#characteristics (from Hatt et al. 2004)
data(urbanwq)
env <- urbanwq[,2:8]
gofs <- all.regs(urbanwq$lec, env, fam = "gaussian",
gof = "Rsqu", print.vars = TRUE)
partition(gofs, pcan = 7, var.names = names(urbanwq[,2,8]))

#hierarchical partitioning of logistic and linear regression
#goodness of fit measures from Chevan and Sutherland (1991)
data(chevan)
partition(chevan$chisq, pcan = 4)
partition(chevan$rsqu, pcan = 4)
```

rand.hp

Randomization test for hierarchical partitioning

Description

Randomizes elements in each column in xcan and recalculates hier.part num.reps times

Usage

```
rand.hp(y, xcan, family = "gaussian", gof = "RMSPE",
num.reps = 100)
```

Arguments

y	a vector containing the dependent variables
xcan	a dataframe containing the n independent variables
family	family argument of glm
gof	Goodness-of-fit measure. Currently "RMSPE", Root-mean-square 'Prediction' error, "NLL", Negative log Likelihood or "Rsqu", R-squared
num.reps	Number of repeated randomizations

Details

This function is a randomization routine for the hier.part function which returns a matrix of I values (the independent contribution towards explained variance in a multivariate dataset) for all values from "num.reps" randomizations. For each randomization, the values in each variable (i.e each column of xcan) are randomized independently, and hier.part is run on the randomized xcan. As well as the randomized I matrix, the function returns a summary table listing the observed I values, the 95th and 99th percentile values of I for the randomized dataset.

Value

a list containing

- Irands** matrix of num.reps + 1 rows of I values for each dependent variable. The first row contains the observed I values and the remaining num.reps rows contains the I values returned for each randomization.
- Iprobs** data.frame of observed I values for each variable, Z-scores for the generated distribution of randomized Is and an indication of statistical significance. Z-scores are calculated as (observed - mean(randomizations))/sd(randomizations), and statistical significance (*) is based on upper 0.95 confidence limit ($Z \geq 1.65$).

Author(s)

Chris Walsh <Chris.Walsh@sci.monash.edu.au>.

References

Hatt, B. E., Fletcher, T. D., Walsh, C. J. and Taylor, S. L. 2004 The influence of urban density and drainage infrastructure on the concentrations and loads of pollutants in small streams. *Environmental Management* **34**, 112–124.

Mac Nally, R. 2000 Regression and model building in conservation biology, biogeography and ecology: the distinction between and reconciliation of 'predictive' and 'explanatory' models. *Biodiversity and Conservation* **9**, 655–671.

Mac Nally, R. 2002 Multiple regression and inference in conservation biology and ecology: further comments on identifying important predictor variables. *Biodiversity and Conservation* **11**, 1397–1401.

Walsh, C. J., Papas, P. J., Crowther, D., Sim, P. T., and Yoo, J. 2004 Stormwater drainage pipes as a threat to a stream-dwelling amphipod of conservation significance, *Austrogammarus australis*, in southeastern Australia. *Biodiversity and Conservation* **13**, 781–793.

See Also

[hier.part](#), [partition](#)

Examples

```
#linear regression of log(electrical conductivity) in streams
#against four independent variables describing catchment
#characteristics (from Hatt et al. 2004)
data(urbanwq)
env <- urbanwq[,2:5]
rand.hp(urbanwq$lec, env, fam = "gaussian", gof = "Rsqu")$Iprobs

#logistic regression of an amphipod species occurrence in
#streams against four independent variables describing
#catchment characteristics (from Walsh et al. 2004)
data(amphipod)
env1 <- amphipod[,2:5]
```

```
rand.hp(amphipod$australis, env1, fam = "binomial",  
gof = "logLik")$Iprobs
```

Index

*Topic **datasets**

hier.part.data, 6

all.regs, 2, 4, 6, 9

amphipod (hier.part.data), 6

chevan (hier.part.data), 6

combos, 3

hier.part, 3, 4, 10

hier.part.data, 6

partition, 3, 6, 7, 9, 10

rand.hp, 3, 6, 9, 9

urbanwq (hier.part.data), 6