

Case study: *L. monocytogenes* in cold-smoked salmon

R. POUILLOT, M.-L. DELIGNETTE-MULLER, M. CORNU

October 11, 2016

The objective of this case study is to assess the risk of invasive listeriosis from consumption of cold-smoked salmon in France. The process of interest lays from the end of the production line in the factory, when the cold-smoked salmon is vacuum-packed, to the consumption.

The data and the model are adapted to illustrate the use of `mc2d`: the results will not and *should not* be interpreted as an assessment of the actual risk of listeriosis from consumption of cold-smoked salmon. Interested readers could refer to [3] and [2] for a complete risk assessment on that issue.

The model will be developed in a first section, without considering variability or uncertainty (deterministic model). Variability will then be introduced in a second section, and a last section will consider variability and a part of the data uncertainty.

1 The Model

In this section, no variability nor uncertainty is considered. We assess the final level of *L. monocytogenes* in the product, the exposure and the risk of invasive listeriosis for an “average” individual of the “healthy” French population¹.

During the logistic, the retail and the home step, a bacterial growth is modeled considering *i*) the fluctuating temperature during the various stages and; *ii*) the bacterial competition with the food flora. We use the models developed and/or used in [3]. The data are adapted from [3] and [1]:

- The DMS model predicts the bacterial growth during a stage of duration d , when the temperature is fluctuating, with an intra-stage average temperature m_T and an intra-stage standard deviation of the temperature s_T . It is written:

$$N_1 = \min \left(N_0 + \frac{\mu_{ref}}{\ln(10)} \times d \times \frac{(s_T^2 + (m_T - T_{min})^2)}{(T_{ref} - T_{min})^2}, N_{max} \right) \quad (1)$$

if $m_T > T_{min}$, with N_1 the \log_{10} concentration of bacteria (\log_{10} (CFU/g)) in the product at the end of the stage, N_0 the \log_{10} concentration of bacteria (\log_{10} (CFU/g)) in the product at the beginning of the stage, μ_{ref} the specific growth rate (day^{-1}) at a reference temperature T_{ref} ($^{\circ}\text{C}$), T_{min} the minimal temperature ($^{\circ}\text{C}$) of growth and N_{max} the maximum achievable concentration in the product (\log_{10} (CFU/g)). If $m_T \leq T_{min}$, $N_1 = N_0$.

- We will use $T_{ref} = 25^{\circ}\text{C}$. We have in this section $N_{max} = 7.27 \log_{10}(\text{CFU/g})$;
- The model for *L. monocytogenes* uses $\mu_{ref,Lm} = 6.2 \text{ day}^{-1}$ and $T_{min,Lm} = -2.9^{\circ}\text{C}$;
- The same model is used for the food flora, with $\mu_{ref,ff} = 4.1 \text{ day}^{-1}$ and $T_{min,ff} = -4.5^{\circ}\text{C}$;
- The growth model for the bacterial competition consider the Jameson effect, i.e. consider that the bacterial growth of *L. monocytogenes* and the growth of the food flora are stopped as soon as one population reaches N_{max} .

¹It makes little sense, but it will help us introducing smoothly the model.

In practice, one will evaluate d_{Lm} and d_{ff} , the time needed for *L. monocytogenes* or the food flora to reach N_{max} , respectively, and model a growth for the given stage during an effective duration of $\min(d, d_{Lm}, d_{ff})$. The time needed to reach N_{max} is evaluated by inverting (1):

$$d_{(N_1=N_{max})} = (N_{max} - N_0) \times \frac{\ln(10)}{\mu_{ref}} \times \frac{(T_{ref} - T_{min})^2}{(s_T^2 + (m_T - T_{min})^2)}$$

The other assumptions are:

- A cold-smoked salmon package is homogeneously contaminated with *L. monocytogenes* at the end of the production at a level of 0.1 CFU/g;
- The food flora level at the end of the production is $10^{2.78}$ CFU/g;
- The time-temperature profile is:
 - 1.1 days at an average temperature of 3.2°C from the factory to the retail (logistic step), with an intra-stage standard deviation of the temperature of 2.1 °C;
 - 4.7 days at an average temperature of 5.5°C at retail with an intra-stage standard deviation of the temperature of 1.0 °C;
 - 4.3 days at an average temperature of 8.2°C in the consumer’s home with an intra-stage standard deviation of the temperature of 2.0 °C;
- An healthy, non elderly, non pregnant individual eats 35g of this product;
- The individual dose-response model for this population is a one hit model

$$\Pr(\text{Illness} | D) = 1 - (1 - r)^D$$

with $r = 4.7 \times 10^{-14}$ for an individual from this healthy sub-population. The populational dose-response that evaluates the mean risk for a population exposed to food where the number of bacteria follows a Poisson distribution of mean parameter D is the exponential dose-response

$$\Pr(\text{Illness} | D) = 1 - \exp(r \times D)$$

The question is “What is the risk for this ‘average’ individual?”. One way to write this model is as following:

```
> Nmax <- 7.3
> murefLm <- 6.2; TminLm <- -2.9
> murefFF <- 4.1; TminFF <- -4.5
> Lm0 <- log10(1); FF0 <- 2.78
> d1 <- 1.1; mT1 <- 3.2; sdT1 <- 2.1
> d2 <- 4.7; mT2 <- 5.5; sdT2 <- 1.0
> d3 <- 4.3; mT3 <- 8.2; sdT3 <- 2.0
> conso <- 35
> r <- 4.7e-14
> modGrowth <- function(duration, mTemp, sdTemp,
+                       NOLm, murefLm, TminLm,
+                       NOFF, murefFF, TminFF,
+                       Nmax, Tref=25) {
+   NOLm <- pmin(NOLm, Nmax)
+   NOFF <- pmin(NOFF, Nmax)
+   dLm <- (Nmax-NOLm) * log(10)/murefLm * (Tref-TminLm)^2 / (sdTemp^2 + (mTemp-TminLm)^2)
+   dLm <- ifelse(mTemp < TminLm & NOLm!=Nmax, Inf, dLm)
+   dFF <- (Nmax-NOFF) * log(10)/murefFF * (Tref-TminFF)^2 / (sdTemp^2 + (mTemp-TminFF)^2)
```

```

+ dFF <- ifelse(mTemp < TminFF & NOFF!=Nmax, Inf, dFF)
+ realDuration <- pmin(duration, dLm , dFF)
+ xLm <- NOLm + (mTemp > TminLm) * murefLm/log(10) *
+       (sdTemp^2 + (mTemp - TminLm)^2) / ((Tref - TminLm)^2) * realDuration
+ xFF <- NOFF + (mTemp > TminFF) * murefFF/log(10) *
+       (sdTemp^2 + (mTemp - TminFF)^2) / ((Tref - TminFF)^2) * realDuration
+ return(list(xLm = xLm, xFF=xFF))}
> x1 <- modGrowth(d1, mT1, sdT1,
+               Lm0, murefLm, TminLm,
+               FF0, murefFF, TminFF,
+               Nmax)
> x2 <- modGrowth(d2, mT2, sdT2,
+               x1$xLm, murefLm, TminLm,
+               x1$xFF, murefFF, TminFF,
+               Nmax)
> x3 <- modGrowth(d3, mT3, sdT3,
+               x2$xLm, murefLm, TminLm,
+               x2$xFF, murefFF, TminFF,
+               Nmax)
> x3

$xLm
[1] 3.21

$xFF
[1] 5.35

> conta <- -10^x3$xLm
> conta

[1] 1637

> expo <- conso * conta
> expo

[1] 57281

> risk <- 1 - (1 - r)^expo
> risk

[1] 2.69e-09

```

`modGrowth` is a convenient function for the growth model. Within this function `dLm` is the time needed for *L. monocytogenes* to reach `Nmax`, `dFF` is the time needed for the food flora to reach `Nmax` and, `realDuration` is the effective time of growth during the stage. Note that:

- this function is “vectorized”, meaning that it can deal with a vector for any of its parameters, returning consequently a vector. This is a strength of R, notably for Monte-Carlo simulations, but it requests a bit of knowledge on the way to code the functions. As an example: `pmin`, a function that takes one or more vectors as arguments and return a single vector giving the “parallel” minima of the vectors is used instead of the more classical function `min` function, that would return the maximum or minimum of *all* the values. Another example is the use of the `ifelse` instead of `if`;
- it is also written to handle *all* specific cases that could occur in the Monte-Carlo simulation, such as $N_0 \geq N_{max}$ or $m_T \leq T_{min}$ or both, for any or both bacterial populations.

`x1`, `x2` and `x3` are the bacterial concentrations at the end of the logistic, the retail and the home step, respectively.

2 Including Variability

We now specify now some variability distributions for some inputs, following [1] and [3]. We first have to call the needed libraries, and define the desired number of iterations:

```
> library(fitdistrplus)
> library(mc2d)
> ndvar(10001)
```

```
[1] 10001
```

2.1 Specifying Variability Distribution

2.1.1 Initial Contamination

For the initial contamination levels in *L. monocytogenes*, we have a set of 62 enumeration data from a representative sample of packages of cold smoked salmon positive in detection: 43 samples have less than 0.2 CFU/g, 7 samples have 0.2 CFU/g, 4 samples have 0.4 CFU/g, 2 samples have 0.6 CFU/g, and the other values are 0.3, 1.0, 1.6, 2.4, 5.4 and 7.0 CFU/g [3]. We will use the `fitdistrplus` package to fit a normal distribution on the \log_{10} of these values, taking into account the censored values. Using the fitted parameters, we model thereafter these initial concentrations in contaminated packages through a normal distribution truncated² on $[-2, \infty)$ \log_{10} (CFU/g).

For the food flora, we use the distribution proposed by [1], $N_{off} \sim N(2.78, 1.14)$.

```
> dataC <- data.frame(
+   left =c(rep(NA,43), rep(.2,7), .3,rep(.4,4),1,1.6,.6,.6,2.4,5.4,7),
+   right=c(rep(0.2,43),rep(.2,7), .3,rep(.4,4),1,1.6,.6,.6,2.4,5.4,7)
+ )
> fit <- fitdistcens(log10(dataC), "norm")
> fit
```

Fitting of the distribution ' norm ' on censored data by maximum likelihood

Parameters:

```
      estimate
mean  -1.117
sd     0.764
```

Fixed parameters:

data frame with 0 columns and 0 rows

```
> LmOV <- mcstoc(rnorm, mean = fit$est["mean"], sd = fit$est["sd"], rtrunc=TRUE, linf=-2)
> FFOV <- mcstoc(rnorm, mean=2.78, sd=1.14)
```

Note that, by default, the type of alea that is modeled is “variability” (`type="V"`).

2.1.2 Growth Parameters

Distributions are derived from [1]:

- N_{max} follows a normal distribution with mean $7.27 \log_{10}$ CFU/g and standard deviation $0.86 \log_{10}$ CFU/g;
- The specific growth rate at the reference temperature of 25°C for *L. monocytogenes* follows a normal distribution with mean 6.24 day^{-1} and standard deviation 0.75 day^{-1} truncated on $[0, \infty)$. The minimal growth temperature follows a normal distribution with mean -2.86°C and standard deviation 1.93°C ;
- The specific growth rate at the reference temperature of 25°C for the food flora follows a normal distribution with mean 4.12 day^{-1} and standard deviation 1.97 day^{-1} truncated on $[0, \infty)$. The minimal growth temperature follows a normal distribution with mean -4.52°C and standard deviation 7.6°C .

²so that at least one CFU is included in one 100g package

Table 1: Time Temperature Profiles

Stage	Mean Temperature (°C)	Intra-Stage Variance of T (°C)	time (days)
logistic	normal(3.2, 2.2) truncated on [-3;25]	$\Gamma(1.16, 4.61)$	Exponential(1.1)
retail	normal(5.5, 2.2) truncated on [-3;25]	$\Gamma(0.65, 2.09)$	Exponential(4.7)
consumer	normal(8.2, 3.8) truncated on [-3; 25]	$\Gamma(0.35, 19.7)$	Exponential(4.3)

```
> NmaxV <- mcstoc(rnorm, mean=7.27, sd = 0.86)
> murefLmV <- mcstoc(rnorm, mean = 6.24, sd = 0.75, rtrunc=TRUE, linf=0)
> TminLmV <- mcstoc(rnorm, mean = -2.86, sd = 1.93)
> murefFFV <- mcstoc(rnorm, mean = 4.12, sd = 1.97, rtrunc=TRUE, linf=0)
> TminFFV <- mcstoc(rnorm, mean = -4.52, sd = 7.66)
```

2.1.3 Time-Temperature Profiles

The time temperature profiles in the three steps are modelled using the distribution provided in the table 1 (adapted from [3] from representative data from France)³. We assume a shelf life of 28 days. A simple way to model this shelf life will be to have $d_1 + d_2 + d_3 \leq 28$ days, with d_1 the duration of the logistic stage, d_2 the duration of the retail stage and d_3 the duration of the consumer stage⁴;

```
> d1V <- mcstoc(rexp, rate = 1/1.1)
> mT1V <- mcstoc(rnorm, mean = 3.2, sd = 2.2, rtrunc = TRUE, linf = -3, lsup = 25)
> sdT1V <- sqrt(mcstoc(rgamma, shape = 1.16, scale=4.61))
> d2V <- mcstoc(rexp, rate = 1/4.7, rtrunc=TRUE, lsup=28-d1V)
> mT2V <- mcstoc(rnorm, mean = 5.5, sd = 2.2, rtrunc = TRUE, linf = -3, lsup = 25)
> sdT2V <- sqrt(mcstoc(rgamma, shape = 0.65, scale=2.09))
> d3V <- mcstoc(rexp, rate = 1/4.3, rtrunc=TRUE, lsup=28-(d1V+d2V))
> mT3V <- mcstoc(rnorm, mean = 8.2, sd = 3.8, rtrunc = TRUE, linf = -3, lsup = 25)
> sdT3V <- sqrt(mcstoc(rgamma, shape = 0.35, scale=19.7))
```

2.1.4 Serving Size

As for the serving size, we consider, from observed data, a discrete empirical distribution with values [3]: $V=\{10, 12, 19, 20, 30, 34, 40, 50, 60, 67.5, 80, 100, 250\}$ grams, observed $F=\{11, 1, 1, 29, 12, 1, 41, 4, 4, 1, 4, 1, 1\}$ time, respectively.

```
> consoV <- mcstoc(rempricalD,
+ values = c(10, 12, 19, 20, 30, 34, 40, 50, 60, 67.5, 80, 100, 250),
+ prob = c(11, 1, 1, 29, 12, 1, 41, 4, 4, 1, 4, 1, 1))
```

2.2 Applying the Model

The model may then be evaluated straightforwardly:

```
> r <- mcdata(4.7e-14, type = "0")
> x1V <- modGrowth(d1V, mT1V, sdT1V,
+ LmOV, murefLmV, TminLmV,
+ FFOV, murefFFV, TminFFV,
+ NmaxV)
> x2V <- modGrowth(d2V, mT2V, sdT2V,
+ x1V$xLm, murefLmV, TminLmV,
```

³ Γ is the Gamma distribution parameterized as $\Gamma(shape, scale)$. The Exponential(x) ditribution is the exponential distribution with mean x .

⁴See the code for a way to model this shelf life using truncated distributions.

```

+           x1V$xFF, murefFFV, TminFFV,
+           NmaxV)
> x3V <- modGrowth(d3V, mT3V, sdT3V,
+           x2V$xLm, murefLmV, TminLmV,
+           x2V$xFF, murefFFV, TminFFV,
+           NmaxV)
> contaV <- 10^-x3V$xLm
> expoV <- consoV * contaV
> riskV <- 1 - exp(-r * expoV )
> Lm1 <- mc(Lm0V, FF0V, NmaxV, murefLmV, TminLmV, murefFFV, TminFFV,
+           d1V, mT1V, sdT1V, d2V, mT2V, sdT2V, d3V, mT3V, sdT3V,
+           consoV, r, contaV, expoV, riskV)
> Lm1

```

	node	mode	nsv	nsu	nva	variate	min	mean	median	max	Nas	type	outm
1	Lm0V	numeric	10001	1	1	1	-2.00e+00	-9.30e-01	-9.88e-01	1.76e+00	0	V	each
2	FF0V	numeric	10001	1	1	1	-1.28e+00	2.78e+00	2.78e+00	6.85e+00	0	V	each
3	NmaxV	numeric	10001	1	1	1	3.97e+00	7.26e+00	7.27e+00	1.06e+01	0	V	each
4	murefLmV	numeric	10001	1	1	1	2.85e+00	6.24e+00	6.25e+00	9.25e+00	0	V	each
5	TminLmV	numeric	10001	1	1	1	-1.01e+01	-2.83e+00	-2.85e+00	3.69e+00	0	V	each
6	murefFFV	numeric	10001	1	1	1	1.31e-02	4.19e+00	4.17e+00	1.13e+01	0	V	each
7	TminFFV	numeric	10001	1	1	1	-3.51e+01	-4.52e+00	-4.46e+00	2.66e+01	0	V	each
8	d1V	numeric	10001	1	1	1	5.36e-05	1.10e+00	7.69e-01	9.69e+00	0	V	each
9	mT1V	numeric	10001	1	1	1	-2.98e+00	3.20e+00	3.15e+00	1.14e+01	0	V	each
10	sdT1V	numeric	10001	1	1	1	3.37e-02	2.08e+00	1.96e+00	6.57e+00	0	V	each
11	d2V	numeric	10001	1	1	1	1.70e-03	4.69e+00	3.29e+00	2.67e+01	0	V	each
12	mT2V	numeric	10001	1	1	1	-2.53e+00	5.53e+00	5.50e+00	1.35e+01	0	V	each
13	sdT2V	numeric	10001	1	1	1	1.60e-03	9.66e-01	8.63e-01	4.76e+00	0	V	each
14	d3V	numeric	10001	1	1	1	3.84e-04	4.09e+00	2.89e+00	2.53e+01	0	V	each
15	mT3V	numeric	10001	1	1	1	-2.98e+00	8.24e+00	8.21e+00	2.26e+01	0	V	each
16	sdT3V	numeric	10001	1	1	1	1.31e-06	1.91e+00	1.41e+00	1.16e+01	0	V	each
17	consoV	numeric	10001	1	1	1	1.00e+01	3.55e+01	4.00e+01	2.50e+02	0	V	each
18	r	numeric	1	1	1	1	4.70e-14	4.70e-14	4.70e-14	4.70e-14	0	0	each
19	contaV	numeric	10001	1	1	1	1.16e-02	4.09e+06	2.22e+01	5.04e+09	0	V	each
20	expoV	numeric	10001	1	1	1	1.70e-01	1.18e+08	6.85e+02	1.01e+11	0	V	each
21	riskV	numeric	10001	1	1	1	7.99e-15	5.55e-06	3.22e-11	4.73e-03	0	V	each

```

> sLm1 <- mc(contaV=Lm1$contaV, expoV=Lm1$expoV, riskV=Lm1$riskV)
> summary(sLm1, probs = c(0, 0.5, 0.75, 0.95, 1))

```

contaV :

	mean	sd	Min	50%	75%	95%	Max	nsv	Na's
NoUnc	4092015	76520808	0.0116	22.2	867	1962680	5.04e+09	10001	0

expoV :

	mean	sd	Min	50%	75%	95%	Max	nsv	Na's
NoUnc	1.18e+08	1.88e+09	0.17	685	27251	60077024	1.01e+11	10001	0

riskV :

	mean	sd	Min	50%	75%	95%	Max	nsv	Na's
NoUnc	5.55e-06	8.82e-05	7.99e-15	3.22e-11	1.28e-09	2.82e-06	0.00473	10001	0

Lm1 is a mc object that contains all the parameters and outputs. We extract some of these outputs in sLm1 to provide a short summary.

2.3 Final Estimate

If 6.5% of cold-smoked salmon package are contaminated, if 49,090,000 Frenchs are part of the “non susceptible” population and if, on average, those people consume some smoked salmon 6.4 times per year, the expected number of cases of listeriosis from consumption of cold smoked salmon in this population is estimated through:

```
> meanRisk <- mcapply(riskV, "var", mean)
> expectedN <- round(0.065 * unmc(meanRisk) * 6.4 * 49090000)
> expectedN
```

```
[1] 113
```

3 Including (a Part of the) Uncertainty

We eventually include both variability and uncertainty in the model. For this example, we will only consider the uncertainty linked to the initial contamination, the growth parameters and the prevalence.

3.1 Specifying Uncertainty

3.1.1 Initial Contamination

The uncertainty surrounding the initial contamination levels of the *L. monocytogenes* will be modeled using a bootstrap procedure, obtained straightforwardly with the help of the `fitdistrplus` package and its `bootdistcens` function. Before this, we define the number of iterations needed in the uncertainty dimension.

```
> ndunc(101)
```

```
[1] 101
```

```
> bootLm0 <- bootdistcens(fit, niter=ndunc())
> MLm0 <- mcdata(bootLm0$est$mean, type="U")
> SLm0 <- mcdata(bootLm0$est$sd, type="U")
> Lm0VU <- mcstoc(rnorm, type="VU", mean=MLm0, sd=SLm0, rtrunc=TRUE, linf=-2)
```

In order to consider uncertainty for the food flora initial contamination, we have, from [1], a set of uncertain hyperparameters, $M_{N_{off}}$ and $\sigma_{N_{off}}$, that are used as parameters for the uncertain *and* variable parameter N_{off} :

$$\begin{aligned} N_{off} &\sim N(M_{N_{off}}, \sigma_{N_{off}}) \\ M_{N_{off}} &\sim N(2.78, 0.265) \\ \ln(\sigma_{N_{off}}) &\sim N(0.114, 0.172) \end{aligned}$$

This hierarchical simulation is written with `mc2d`:

```
> MLmOFF <- mcstoc(rnorm, type="U", mean=2.78, sd=0.265)
> SLmOFF <- mcstoc(rlnorm, type="U", meanlog=0.114, sdlog=0.172)
> FFOVU <- mcstoc(rnorm, type="VU", mean=MLmOFF, sd=SLmOFF)
```

3.1.2 Growth Parameters

The uncertainty around $\mu_{ref,Lm}$, $T_{min,Lm}$, $\mu_{ref,ff}$, $T_{min,ff}$ and N_{max} are modeled similarly through the specification of hyperparameters [1]⁵:

$$\begin{aligned}\mu_{ref,Lm} &\sim N(M_{\mu_{ref,Lm}}, \sigma_{\mu_{ref,Lm}}) \\ M_{\mu_{ref,Lm}} &\sim \Gamma(shape : 69.7, scale : 0.0896) \\ \ln(\sigma_{\mu_{ref,Lm}}) &\sim N(1.03, 0.191)\end{aligned}$$

$$\begin{aligned}T_{min,Lm} &\sim N(M_{T_{min,Lm}}, \sigma_{T_{min,Lm}}) \\ M_{T_{min,Lm}} &\sim N(-2.86, 0.459) \\ \ln(\sigma_{T_{min,Lm}}) &\sim N(0.638, 0.208)\end{aligned}$$

$$\begin{aligned}\mu_{ref,ff} &\sim N(M_{\mu_{ref,ff}}, \sigma_{\mu_{ref,ff}}) \\ M_{\mu_{ref,ff}} &\sim \Gamma(shape : 32.5, scale : 0.127) \\ \ln(\sigma_{\mu_{ref,ff}}) &\sim N(-0.656, 0.221)\end{aligned}$$

$$\begin{aligned}T_{min,ff} &\sim N(M_{T_{min,ff}}, \sigma_{T_{min,ff}}) \\ M_{T_{min,ff}} &\sim N(-4.52, 1.23) \\ \ln(\sigma_{T_{min,ff}}) &\sim N(2.00, 0.257)\end{aligned}$$

$$\begin{aligned}N_{max} &\sim N(M_{N_{max}}, \sigma_{N_{max}}) \\ M_{N_{max}} &\sim N(7.27, 0.276) \\ \ln(\sigma_{N_{max}}) &\sim N(-0.172, 0.218)\end{aligned}$$

with $\mu_{ref} > 0$ and $T_{min} < 25$. We simply translated the preceding distributions:

```
> MmurefLm <- mcstoc(rgamma, type="U", shape=69.7, scale=0.0896)
> SmurefLm <- mcstoc(rlnorm, type="U", meanlog = 1.03, sdlog = 0.191)
> murefLmVU <- mcstoc(rnorm, type="VU", mean=MmurefLm, sd=SmurefLm, rtrunc=TRUE, linf=0)
> MTminLm <- mcstoc(rnorm, type="U", mean=-2.86, sd=0.459)
> STminLm <- mcstoc(rlnorm, type="U", meanlog = 0.638, sdlog = 0.208)
> TminLmVU <- mcstoc(rnorm, type="VU", mean = MTminLm, sd = STminLm, rtrunc=TRUE, lsup=25)
> MmurefFF <- mcstoc(rgamma, type="U", shape=32.5, scale=.127)
> SmurefFF <- mcstoc(rlnorm, type="U", meanlog = -.656, sdlog = 0.221)
> murefFFVU <- mcstoc(rnorm, type="VU", mean=MmurefFF, sd=SmurefFF, rtrunc=TRUE, linf=0)
> MTminFF <- mcstoc(rnorm, type="U", mean=-4.52, sd=1.23)
> STminFF <- mcstoc(rlnorm, type="U", meanlog = 2.00, sdlog = 0.257)
> TminFFVU <- mcstoc(rnorm, type="VU", mean = MTminFF, sd = STminFF, rtrunc=TRUE, lsup=25)
> MNmax <- mcstoc(rnorm, type="U", mean=7.27, sd=0.276)
> SNmax <- mcstoc(rlnorm, type="U", meanlog = -0.172, sdlog = 0.218)
> NmaxVU <- mcstoc(rnorm, type="VU", mean = MNmax, sd = SNmax)
```

3.1.3 Prevalence

The prevalence level of contaminated cold-smoked salmon packages (6.5%) was estimated from 41 positive packages out of 626 tested [3]. We assume a sensitivity and a specificity of the method of 100%. We model the data

⁵Note that there was a typo in [1] that lead to an error in [3]: the standard-error for $\ln(\sigma_{\mu_{ref,Lm}})$ is 1.03 and not -1.03 as written in [1]. We will use here the correct value.

uncertainty around the true prevalence of contaminated package using a bayesian reasoning, with a Beta(1, 1) distribution as a prior. The number of expected cases may be estimated using:

```
> prevU <- mcstoc(rbeta,type="U", shape1=41+1, shape2=626-41+1)
```

3.2 Applying the Model

Applying the model is just a copy-paste from the previous version (+ we change the name of the parameters).

```
> x1VU <- modGrowth(d1V, mT1V, sdT1V,
+                 LmOVU, murefLmVU, TminLmVU,
+                 FFOVU, murefFFVU, TminFFVU,
+                 NmaxVU)
> x2VU <- modGrowth(d2V, mT2V, sdT2V,
+                 x1VU$xLm, murefLmVU, TminLmVU,
+                 x1VU$xFF, murefFFVU, TminFFVU,
+                 NmaxVU)
> x3VU <- modGrowth(d3V, mT3V, sdT3V,
+                 x2VU$xLm, murefLmVU, TminLmVU,
+                 x2VU$xFF, murefFFVU, TminFFVU,
+                 NmaxVU)
> contaVU <- 10^x3VU$xLm
> expoVU <- consoV * contaVU
> riskVU <- 1 - exp(-r * expoVU)
> Lm2 <- mc(LmOVU, FFOVU, NmaxVU, murefLmVU, TminLmVU, murefFFVU, TminFFVU,
+          d1V, mT1V, sdT1V, d2V, mT2V, sdT2V, d3V, mT3V, sdT3V,
+          consoV, r, contaVU, expoVU, riskVU)
> Lm2
```

	node	mode	nsv	nsu	nva	variate	min	mean	median	max	Nas	type	outm
1	LmOVU	numeric	10001	101	1	1	-2.00e+00	-9.37e-01	-9.93e-01	3.68e+00	0	VU	each
2	FFOVU	numeric	10001	101	1	1	-4.82e+00	2.76e+00	2.76e+00	9.58e+00	0	VU	each
3	NmaxVU	numeric	10001	101	1	1	2.11e+00	7.28e+00	7.28e+00	1.27e+01	0	VU	each
4	murefLmVU	numeric	10001	101	1	1	1.00e-04	6.45e+00	6.36e+00	2.43e+01	0	VU	each
5	TminLmVU	numeric	10001	101	1	1	-1.44e+01	-2.83e+00	-2.84e+00	8.70e+00	0	VU	each
6	murefFFVU	numeric	10001	101	1	1	4.81e-03	4.21e+00	4.21e+00	8.24e+00	0	VU	each
7	TminFFVU	numeric	10001	101	1	1	-5.90e+01	-4.39e+00	-4.35e+00	2.50e+01	0	VU	each
8	d1V	numeric	10001	1	1	1	5.36e-05	1.10e+00	7.69e-01	9.69e+00	0	V	each
9	mT1V	numeric	10001	1	1	1	-2.98e+00	3.20e+00	3.15e+00	1.14e+01	0	V	each
10	sdT1V	numeric	10001	1	1	1	3.37e-02	2.08e+00	1.96e+00	6.57e+00	0	V	each
11	d2V	numeric	10001	1	1	1	1.70e-03	4.69e+00	3.29e+00	2.67e+01	0	V	each
12	mT2V	numeric	10001	1	1	1	-2.53e+00	5.53e+00	5.50e+00	1.35e+01	0	V	each
13	sdT2V	numeric	10001	1	1	1	1.60e-03	9.66e-01	8.63e-01	4.76e+00	0	V	each
14	d3V	numeric	10001	1	1	1	3.84e-04	4.09e+00	2.89e+00	2.53e+01	0	V	each
15	mT3V	numeric	10001	1	1	1	-2.98e+00	8.24e+00	8.21e+00	2.26e+01	0	V	each
16	sdT3V	numeric	10001	1	1	1	1.31e-06	1.91e+00	1.41e+00	1.16e+01	0	V	each
17	consoV	numeric	10001	1	1	1	1.00e+01	3.55e+01	4.00e+01	2.50e+02	0	V	each
18	r	numeric	1	1	1	1	4.70e-14	4.70e-14	4.70e-14	4.70e-14	0	0	each
19	contaVU	numeric	10001	101	1	1	1.01e-02	1.24e+07	1.71e+01	5.74e+11	0	VU	each
20	expoVU	numeric	10001	101	1	1	1.01e-01	4.53e+08	5.22e+02	2.87e+13	0	VU	each
21	riskVU	numeric	10001	101	1	1	4.77e-15	2.03e-05	2.46e-11	7.41e-01	0	VU	each

```
> sLm2 <- mc(contaVU=Lm2$contaVU, expoVU=Lm2$expoVU, riskVU=Lm2$riskVU)
> summary(sLm2, probs = c(0, 0.5, 0.75, 0.95, 1))
```

```

contaVU :
      mean      sd   Min  50%  75%    95%    Max  nsv Na's
median 7587743 1.10e+08 0.0112 15.41  901  4433567 6.87e+09 10001  0
mean   12361541 3.16e+08 0.0116 24.22 2743  9091352 2.52e+10 10001  0
2.5%   1118745 1.05e+07 0.0101  6.18  198   367659 4.77e+08 10001  0
97.5%  48245013 1.65e+09 0.0142 96.93 18853 37453053 1.54e+11 10001  0

```

```

expoVU :
      mean      sd   Min  50%  75%    95%    Max  nsv Na's
median 2.57e+08 4.89e+09 0.149  483 27579 1.35e+08 3.30e+11 10001  0
mean   4.53e+08 1.28e+10 0.160  731 84501 2.83e+08 1.04e+12 10001  0
2.5%   4.15e+07 4.72e+08 0.105  189  6216 1.17e+07 2.27e+10 10001  0
97.5%  1.96e+09 6.28e+10 0.257 2871 581399 1.14e+09 5.63e+12 10001  0

```

```

riskVU :
      mean      sd   Min  50%  75%    95%    Max  nsv Na's
median 1.20e-05 2.28e-04 6.99e-15 2.27e-11 1.30e-09 6.34e-06 0.01539 10001  0
mean   2.03e-05 5.19e-04 7.50e-15 3.43e-11 3.97e-09 1.33e-05 0.04066 10001  0
2.5%   1.95e-06 2.22e-05 4.94e-15 8.88e-12 2.92e-10 5.52e-07 0.00106 10001  0
97.5%  8.82e-05 2.65e-03 1.21e-14 1.35e-10 2.73e-08 5.34e-05 0.23139 10001  0

```

The summary provides the estimate of the mean, the standard deviation, the minimum, the median ... and a 95% credible interval. The estimate is the median of the 101 values obtained in the uncertainty dimension. The credible interval lays between the 2.5th and the 97.5th percentiles obtained in the uncertainty dimension.

3.3 Final Estimate

The uncertainty around the number of expected cases is estimated using:

```

> meanRiskU <- mcapply(riskVU, "var", mean)
> expectedNU <- round(prevU * meanRiskU * 6.4 * 49090000)
> summary(expectedNU)

```

```

node :
      NoVar
median  267
mean    423
2.5%    38
97.5%  1954

```

This is an estimate of the uncertainty around the number of cases linked to the uncertainty around the initial contamination, the bacterial growth parameter and the sampling uncertainty for positive packages. A lot of other uncertainties exist but are not considered here, notably the uncertainty around the dose-response model and parameters. See [3, 2] for a complete analysis. The study of the model through a Tornado chart in the variability dimension leads to the Figure 1. It suggests a big impact of the growth rate of *L. monocytogenes*, of the storage duration during the consumer step, and of the initial level of *L. monocytogenes*. The Tornado chart in the uncertainty dimension leads to the Figure 2 and suggests the impact of the uncertainty around N_{max} on the mean risk, and thus the expected number of cases.

```

> torn <- tornado(Lm2)
> torn

```

```

Spearman's rho statistic
Output: riskVU
$riskVU

```

```

      Lm0VU  FF0VU NmaxVU murefLmVU TminLmVU murefFFVU TminFFVU  d1V  mT1V  sdT1V  d2V
median 0.303 -0.0823 0.0711 0.465 -0.237 -0.02929 0.1184 0.0447 0.0337 0.00419 0.277
mean 0.298 -0.0888 0.0751 0.457 -0.238 -0.03060 0.1257 0.0440 0.0334 0.00361 0.277
2.5% 0.202 -0.1567 0.0293 0.347 -0.342 -0.05684 0.0444 0.0245 0.0197 -0.01155 0.227
97.5% 0.380 -0.0340 0.1382 0.548 -0.159 -0.00703 0.2246 0.0608 0.0488 0.01783 0.328

      mT2V  sdT2V  d3V  mT3V  sdT3V  consoV  contaVU  expoVU
median 0.158 0.00741 0.406 0.260 0.0291 0.125 0.991 1
mean 0.158 0.00714 0.407 0.259 0.0296 0.125 0.990 1
2.5% 0.132 -0.00798 0.330 0.219 0.0165 0.105 0.986 1
97.5% 0.187 0.02019 0.479 0.309 0.0418 0.148 0.994 1

```

```

> tornunc <- tornadounc(Lm2, quant=.975)
> tornunc

```

```

Tornado on uncertainty
Spearman's rho statistic
Output: riskVU
$riskVU

```

```

      mean Lm0VU sd Lm0VU 97.5% Lm0VU mean FF0VU sd FF0VU 97.5% FF0VU mean NmaxVU sd NmaxVU
mean riskVU 0.155 0.04478 0.0549 -0.0904 -0.170 -0.195 0.656 0.720
sd riskVU 0.161 -0.00829 0.0139 -0.0186 -0.179 -0.152 0.509 0.796
97.5% riskVU 0.186 0.17879 0.1750 -0.2009 -0.120 -0.228 0.735 0.242

      97.5% NmaxVU mean murefLmVU sd murefLmVU 97.5% murefLmVU mean TminLmVU sd TminLmVU
mean riskVU 0.912 0.409 0.286 0.439 0.0176 -0.0524
sd riskVU 0.900 0.280 0.179 0.296 0.0736 -0.1391
97.5% riskVU 0.552 0.639 0.428 0.639 -0.0901 0.0917

      97.5% TminLmVU mean murefFFVU sd murefFFVU 97.5% murefFFVU mean TminFFVU sd TminFFVU
mean riskVU -0.0305 -0.1796 0.0659 -0.1505 0.159 0.003891
sd riskVU -0.0806 -0.0773 0.0337 -0.0649 0.140 0.000711
97.5% riskVU 0.0446 -0.3025 0.0494 -0.2760 0.173 0.025172

      97.5% TminFFVU mean contaVU sd contaVU 97.5% contaVU mean expoVU sd expoVU
mean riskVU 0.0269 0.994 0.931 0.780 1.000 0.936
sd riskVU 0.0111 0.928 0.967 0.571 0.938 1.000
97.5% riskVU 0.0569 0.769 0.562 0.997 0.772 0.567

      97.5% expoVU
mean riskVU 0.776
sd riskVU 0.568
97.5% riskVU 1.000

```

```

> plot(torn)
> plot(tornunc, stat="mean risk")

```

As a conclusion, this example illustrates how predictive growth models may be implemented within mc2d...

References

- [1] M.-L. Delignette-Muller, M. Cornu, R. Pouillot, and J.-B. Denis. Use of bayesian modelling in risk assessment: application to growth of *Listeria monocytogenes* and food flora in cold-smoked salmon. *International Journal of Food Microbiology*, 106(2):195–208, 2006.
- [2] R. Pouillot, V. Goulet, M. L. Delignette-Muller, A. Mahe, and M. Cornu. Quantitative risk assessment of listeria monocytogenes in french cold-salmon : Ii. risk characterization. *Risk Analysis*, 29(6):806–819, 2009.

Figure 1: Tornado chart for the *L. monocytogenes* example (Variability).





Figure 2: Tornado chart for the *L. monocytogenes* example (Uncertainty).



- [3] R. Pouillot, N. Miconnet, A.-L. Afchain, M.-L. Delignette-Muller, A. Beaufort, L. Rosso, J.-B. Denis, and M. Cornu. Quantitative risk assessment of listeria monocytogenes in french cold-salmon : I. quantitative exposure assessment. *Risk Analysis*, 27(3):683–700, 2007.