

# Package ‘EcoHydRology’

September 24, 2018

**Version** 0.4.12.1

**Title** A Community Modeling Foundation for Eco-Hydrology

**Author** Fuka DR, Walter MT, Archibald JA, Steenhuis TS, and Easton ZM

**Maintainer** Daniel Fuka <drf28@cornell.edu>

**Depends** R (>= 2.10), operators, topmodel, DEoptim, XML

**Description** Provides a flexible foundation for scientists, engineers, and policy makers to base teaching exercises as well as for more applied use to model complex eco-hydrological interactions.

**License** GPL-2

**Repository** CRAN

**Date/Publication** 2018-09-24 12:52:33 UTC

**KeepSource** TRUE

**NeedsCompilation** no

## R topics documented:

EcoHydRology-package . . . . .	2
alter_files . . . . .	3
AtmosphericEmissivity . . . . .	4
BaseflowSeparation . . . . .	5
build_gsod_forcing_data . . . . .	6
calib_swat_ex . . . . .	7
change_params . . . . .	8
ConvertFlowUnits . . . . .	9
declination . . . . .	10
EnvirEnergy . . . . .	11
EstCloudiness . . . . .	12
EvapHeat . . . . .	13
get_cfsr_latlon . . . . .	14
get_gsod_stn . . . . .	15
get_usgs_gage . . . . .	17
GroundHeat . . . . .	18

GSOD_history . . . . .	19
hydrograph . . . . .	20
Longwave . . . . .	21
Lumped_VSA_model . . . . .	22
NetRad . . . . .	24
OwascoInlet . . . . .	25
PET_fromTemp . . . . .	26
PotentialSolar . . . . .	27
PotSolarInst . . . . .	28
PTpet . . . . .	29
RainHeat . . . . .	29
RunoffBreakdown . . . . .	30
SatVaporDensity . . . . .	30
SatVaporPressure . . . . .	31
SatVapPresSlope . . . . .	32
SensibleHeat . . . . .	32
setup_swatcal . . . . .	33
slopefactor . . . . .	34
SnowMelt . . . . .	35
SoilStorage . . . . .	36
Solar . . . . .	37
solarangle . . . . .	38
SWAT2005 . . . . .	38
swat_general . . . . .	39
swat_objective_function . . . . .	44
swat_objective_function_rch . . . . .	45
testSWAT2005 . . . . .	45
transmissivity . . . . .	46
<b>Index</b>	<b>48</b>

---

EcoHydRology-package    *A community modeling foundation for Eco-Hydrology.*

---

## Description

This package provides a flexible foundation for scientists, engineers, and policy makers to base teaching exercises as well as for more applied use to model complex eco-hydrological interactions.

## Details

Package:	EcoHydRology
Type:	Package
Version:	0.4.12
Date:	2013-09-27
License:	GPL-2
LazyLoad:	yes

**Included Energy Budget Functions:**

AtmosphericEmissivity

EnvirEnergy

EstCloudiness

EvapHeat

GroundHeat

Longwave

NetRad

PotentialSolar

RainHeat

SatVaporDensity

SensibleHeat

Solar

declination

slopefactor

solarangle

solaraspect

transmissivity

**Included Data Handeling Functions:**

get\_gsod\_stn

build\_gsod\_forcing\_data

**Included data:**

GSOD\_history

**Author(s)**

Fuka DR, Walter MT, Archibald JA, Steenhuis, TS, and Easton ZM

Maintainer: Daniel Fuka &lt;drf28@cornell.edu&gt;

---

`alter_files`*A function to alter files called from swat\_objective\_function*

---

**Description**

A function to alter files called from swat\_objective\_function

**Usage**`alter_files(change_params)`**Arguments**`change_params`

**Author(s)**

Daniel R. Fuka

---

AtmosphericEmissivity *Atmospheric Emissivity*

---

**Description**

The emissivity of the atmosphere [-]

**Usage**

AtmosphericEmissivity(airtemp, cloudiness, vp=NULL, opt="linear")

**Arguments**

airtemp	Air temperature: air temperature [C]
cloudiness	Cloudiness: fraction of the sky covered in clouds [-]
vp	Vapor Pressure : [kPa]
opt	option: either "linear" for a linear relationship between clear sky emissivity and temperature, or "Brutsaert" to use Brutsaert's(1975) clear sky emissivity formulation - this requires input for vapor pressure

**Value**

The emissivity of the atmosphere [-]

**Author(s)**

Fuka, D.R., Walter, M.T., Archibald, J.A.

**References**

Campbell, G. S., Norman, J.M., 1998. An Introduction to Environmental Biophysics, second ed., Springer, New York, p. 286.

Brutsaert, W. (1975) On a Derivable Formula for Long-Wave Radiation From Clear Skies. Water Resources Research, 11(5) 742-744

**Examples**

```
temp=15
clouds=.5
AtmEm=AtmosphericEmissivity(temp,clouds)
print(AtmEm)
```

---

BaseflowSeparation      *Baseflow Separation*

---

### Description

This function reads a streamflow dataset and produces a baseflow dataset. It can be run using 1, 2 or 3 passes

### Usage

```
BaseflowSeparation(streamflow, filter_parameter = 0.925, passes = 3)
```

### Arguments

streamflow	A vector containing streamflow values
filter_parameter	The value recommended by Nathan and McMahon (1990) is 0.925, however, the user might want to play with this value (0.9-0.95)
passes	The number of times you want the filter to pass over your data. 1-3

### Value

This will return a 2 column data frame with nrow = length of input streamflow data. The first column contains baseflow, while the second contains quickflow, both in the same units as the input.

### Author(s)

Josephine Archibald

### References

Lyne, V. D. and M. Hollick (1979). Stochastic time-variable rainfall-runoff modelling. Hydrology and Water Resources Symposium, Perth, Institution of Engineers, Australia.

Nathan, R. J. and T. A. McMahon (1990). "Evaluation of automated techniques for base flow and recession analysis." Water Resources Research 26(7): 1465-1473.

### Examples

```
##### Look at a dataset for Owasco Lake in NY:
data(OwascoInlet)
summary(OwascoInlet)

## Get an approximation for baseflow using a 3 pass filter:
bfs<-BaseflowSeparation(OwascoInlet$Streamflow_m3s, passes=3)

## You can check out how this looks with the hydrograph function:
hydrograph(input=OwascoInlet,streamflow2=bfs[,1])
```

---

 build\_gsod\_forcing\_data

*Parsing gzipped GSOD dataset.*


---

### Description

Parsing of the gzipped GSOD forcing data as returned from `get_gsod_stn()`

### Usage

```
build_gsod_forcing_data()
```

### Author(s)

Daniel R. Fuka

### Examples

```
#
# After running get_gsod_stn() with addis as a temp directory then:
#

## Not run:
tmp_ppt<-build_gsod_forcing_data()
addis

## End(Not run)
## The function is currently defined as
function(){
  tmpdir=readline("Please enter a temp directory where you stored the
*.op.gz datafiles? \n")
  files=dir(tmpdir,"*.op.gz",full.names=T)
  start_year=min(substr(files,nchar(files)-9,nchar(files)-6))
  end_year=max(substr(files,nchar(files)-9,nchar(files)-6))
  alldates=data.frame(fdate=seq(from=as.Date(paste(start_year,"-01-01",
sep="")), to=as.Date(paste(end_year,"-12-31",sep="")), by=1))

  stn=matrix()
  tmin=matrix()
  tmax=matrix()
  ppt=matrix()
  fdate=matrix()
  for (tmpfile in files){
    #
    # There is more data in this dataset we can extract later as we need it.
    #
    #
    tmpstring<-grep("MAX",readLines( gzfile(tmpfile)),value=TRUE,invert=TRUE)
    stn<-c(stn,as.numeric(as.character(substring(tmpstring,1,5))))
    tmax<-c(tmax,as.numeric(as.character(substring(tmpstring,103,108))))
```

```

    tmin<-c(tmin,as.numeric(as.character(substring(tmpstring,111,116))))
    ppt<-c(ppt,as.numeric(as.character(substring(tmpstring,119,123))))
    fdate<-c(fdate,as.Date(yearmoda<-substring(tmpstring,15,22),
"%Y%m%d"))
  }

  stn<-as.numeric(stn)
  ppt<-as.numeric(ppt)
  tmax<-as.numeric(tmax)
  tmin<-as.numeric(tmin)
  fdate<-as.Date(as.numeric(fdate), origin="1970-01-01")
  forcing=data.frame(stn=stn,ppt=ppt,tmax=tmax,tmin=tmin,
  fdate=as.Date(fdate))
  forcing=na.omit(forcing)
  forcing=merge(alldates,forcing,all=TRUE)

  forcing$ppt_mm <- forcing$ppt*25.4
  forcing$tmax_C <- (forcing$tmax-32) * 5/9
  forcing$tmin_C <- (forcing$tmin-32) * 5/9
  forcing$tavg_C <-(forcing$tmin_C+forcing$tmax_C)/2
  forcing$ppt_mm[forcing$ppt_mm > 999]=0.0
  return(forcing)
}

```

---

calib\_swat\_ex

*A function demonstrate an example series of steps to calibrate a reach in the SWAT2005 watershed model.*

---

## Description

A function demonstrate an example series of steps to calibrate a reach in the SWAT2005 watershed model. This should act only as a simple demonstration, and should be modified to individuals use.

## Usage

```
calib_swat_ex(flowgage, rch = 3)
```

## Arguments

flowgage	A list in the format that would be returned from the <code>get_usgs_gage</code>
rch	The reach in the output.rch file you wish to calibrate against.

## Details

This should act only as a simple demonstration, and should be modified to individuals use.

## Value

List (of length 2) containing the results of the internally called `DEoptim` function. See `DEoptim` for more information.

**Author(s)**

Daniel R. Fuka

**Examples**

```

## Not run:
#
install.packages("SWATmodel")
#
# Load library, test SWAT numerics, define a flow gage id for USGS gage, or build similar list
# of data as presented in ?get_usgs_gage
#
library(SWATmodel)
testSWAT2005()
flowgage_id="04216500"
flowgage=get_usgs_gage(flowgage_id)
#
# Obtain basic set of historical forcing data from TAMU/Cornell/IWMI CFSR repository
#
hist_wx=get_cfsr_latlon(flowgage$declat,flowgage$declon)
#
# Build a generic 9 HRU SWAT initialization, which builds and changes into directory
# named as flowgage_id above.
#
build_swat_basic(dirname=flowgage_id,iyr="1979",nbyr=6,flowgage$area,
flowgage$elev,flowgage$declat,flowgage$declon,hist_wx=hist_wx)
#
# Simple calibration based on the outflow from RCH 3 of the simple SWAT initialization
#
calib_swat_results=calib_swat_ex(flowgage,rch=3)
#
# Graph output as hydro graph and pred/obs graphics
plot(calib_swat_results$flowdata,flowgage$flowdata)

## End(Not run)

```

---

change\_params

*Example dataframe of parameters, ranges, and search strings for optimizing swat2005*


---

**Description**

Example dataframe of parameters, ranges, and search strings for optimizing swat2005

**Usage**

```
data(change_params)
```



**Format**

A data frame with 15 observations on the following 11 variables.

filetype a factor with levels 0\*.gw 0\*.mgt 0\*.sol basins.bsn

parameter a factor with levels ALPHA\_BF Ave\ AW Bulk Density CN2 Depth ESCO GW\_DELAY  
GWQMN Ksat\ \ (est\.\) SFTMP SMFMN SMFMX SMTMP SURLAG TIMP

alter\_type a factor with levels new percent

min a numeric vector

max a numeric vector

current a numeric vector

multi a logical vector

frformat a factor with levels A27,10f12

fwformat a factor with levels F16.4 F20

**Examples**

```
data(change_params)
## maybe str(change_params) ; plot(change_params) ...
```

---

ConvertFlowUnits      *Unit conversion for flow rates*

---

**Description**

Converts volumetric flow (cfs, cms, cmd) to depth flow rate over a watershed (mm/d). Or, it converts a depth flow rate to volumetric (mm/d to cfs)

**Usage**

```
ConvertFlowUnits(cfs = NULL, cmd=NULL, cms = NULL, WA, mmd = NULL, AREAunits = "mi2")
```

**Arguments**

cfs	Input flow in cubic feet per second
cmd	Input flow in cubic meters per day
cms	Input flow in cubic meters per second
WA	Watershed area. Can be entered in square miles (default), or square km (in this case, change the AREAunits to "km2")
mmd	Input flow in mm/d
AREAunits	Units of the watershed area ("mi2" or "km2")



**Author(s)**

M. Todd Walter

**Examples**

```
##---- Should be DIRECTLY executable !! ----
##-- ==> Define data, use random,
##--or do help(data=index) for the standard data sets.

## The function is currently defined as
function(Jday){
# solar declination [rad]
#
#Jday: Julian date or day of the year [day]
#
return(0.4102*sin(pi*(Jday-80)/180))
}
```

---

 EnvirEnergy

*The Surface Net Energy Budget*


---

**Description**

Net Energy=S+La-Lt+S+E+R+G, where S is the net incident solar radiation, La is the atmospheric long wave radiation, Lt is the terrestrial long wave radiation, H is the sensible heat exchange, E is the energy flux associated with the latent heats of vaporization and condensation, G is ground heat conduction, P is heat added by precipitation

**Usage**

```
EnvirEnergy(lat, Jday, Tx, Tn, wind, rain, relativehumidity,
  albedo=0.2, cloudiness=NULL, forest=0, slope=0, aspect=0,
  surftemp=(Tx+Tn)/2, surfemissivity=0.97)
```

**Arguments**

lat	latitdue [rad]
Jday	Julian day / day of the year, 1-365 [day]
Tx	Maximum daily temperature [C]
Tn	Minimum daily temperature [C]
wind	Average daily windspeed [m/s]
rain	precipitation [mm/day]
relativehumidity	Relative humidity; if negative, air vapor density will be approximated [-]
albedo	#albedo: surface albedo or reflectivity [-]

cloudiness	Fraction of the sky covered in clouds,if no value provided, cloudiness will be approximated [-]
forest	Forest or vegetation cover [-]
slope	Slope of the ground [rad]
aspect	Ground aspect [rad from north]
surftemp	surface temperature [C]
surfemissivity	surface emissivity: [-]

**Value**

NetEnergy      Surface net energy

**Author(s)**

Walter, M.T.,Fuka, D.R. Maintainer: Daniel Fuka <drf28@cornell.edu>

**Examples**

```
## The function is currently defined as
function(lat,Jday,Tx,Tn,wind,relativehumidity,cloudiness,albedo,forest,
slope,aspect,surftemp,surfemissivity){
if(cloudiness<0){cloudiness<-EstCloudiness(lat,Jday,Tx,Tn)}

airtemp<-(Tx+Tn)/2 #average daily air temperature [C]

return(Solar(lat,Jday,Tx,Tn,albedo,forest,slope,aspect)+
Longwave(AtmosphericEmissivity(airtemp,cloudiness),airtemp)-
Longwave(surfemissivity,surftemp)+SensibleHeat(surftemp,airtemp,wind)+
EvapHeat(surftemp,airtemp,relativehumidity,Tn,wind)+
RainHeat(airtemp,rain)+GroundHeat())
}
```

---

EstCloudiness	<i>Estimated Cloudiness</i>
---------------	-----------------------------

---

**Description**

Estimates the cloudiness of the atmosphere by scaling to atmospheric transmissivity

**Usage**

```
EstCloudiness(Tx=(-999), Tn=(-999), trans=NULL, transMin = 0.15,
transMax = 0.75, opt = "linear")
```

**Arguments**

Tx	maximum daily temperature [C]
Tn	minimum daily temperature [C]
trans	transmissivity of the atmosphere (value between 0-1)
transMin	Transmissivities equal and below this value will return a cloudiness value of 1 (for linear approximation)
transMax	Transmissivities equal and above this value will return a cloudiness value of 0 (for linear approximation)
opt	Currently there are two options: "linear" (Campbell 1985) and "Black" (1956)

**Author(s)**

M. Todd Walter, Josephine Archibald

**References**

Campbell G.S. (1985) Soil Physics with Basic: Transport Models for Soil Plant Systems. Elsevier, New York

Black, J.N. (1956) The Distribution of Solar Radiation over the Earth's Surface. Theoretical and Applied Climatology 7:2 165-189

**Examples**

```
EstCloudiness(trans=0.2)
EstCloudiness(20, 12, opt="Black")
```

---

EvapHeat	<i>Evaporative heat exchange between a wet surface and the surrounding air</i>
----------	--

---

**Description**

Evaporative heat exchange between a surface and the surrounding air [kJ m<sup>-2</sup> d<sup>-1</sup>]. This function is only intended for wet surfaces, i.e., it assumes the vapor density at the surface is the saturation vapor density

**Usage**

```
EvapHeat(surftemp, airtemp, relativehumidity=NULL, Tn=NULL, wind=2)
```

**Arguments**

surftemp	surface temperature [C]
airtemp	average daily air temperature [C]
relativehumidity	relative humidity, 0-1 [-]
Tn	minimum daily air temperature, assumed to be dew point temperature if relativehumidity unknown [C]
wind	average daily windspeed [m/s]

**Note**

This function will use the relative humidity to estimate air vapor density if the value passed is between 0-1. Otherwise, it will assume the minimum daily air temperature is approximately the dew point temp.

**Author(s)**

M. Todd Walter

**References**

Campbell, Gaylon S. An Introduction to Environmental Biophysics. New York: Springer-Verlag, 1977.

**Examples**

```
EvapHeat(surftemp=15, airtemp=5, relativehumidity=0.7)
```

---

get_cfsr_latlon	<i>This is a function to grab daily summaries of the CFSR from the drfuka.org service</i>
-----------------	---

---

**Description**

This is a function to grab daily summaries of the CFSR from the drfuka.org service

**Usage**

```
get_cfsr_latlon(declat, declon, emailaddr, timeoff = 0, interppow = 2)
```

**Arguments**

```
declat
declon
emailaddr
timeoff
interppow
```

**Author(s)**

Daniel R Fuka

**Examples**

## Not run:

```
hist_wx=get_cfsr_latlon(45,-72,"dan@dan.com",timeoff=0,interppow=2)
plot(hist_wx$TMX)
```

## End(Not run)

## The function is currently defined as

```
function (declat, declon, emailaddr, timeoff = 0, interppow = 2)
```

```
{
```

```
  library(XML)
```

```
  options(timeout = 120)
```

```
  url = paste("http://www.cfsr.tamu-cornell.drfuka.org/swat-cfsr-v02.pl?lat=",
```

```
    declat, "&lon=", declon, "&timeoff=", timeoff, "&interppow=",
```

```
    interppow, "&.submit=Submit", sep = "")
```

```
  hist_wx = readHTMLTable(url, which = 1, header = T, colClasses = c("character",
```

```
    "numeric", "numeric", "numeric", "numeric", "numeric",
```

```
    "numeric", "numeric", "numeric"))
```

```
  hist_wx$DATE = as.Date(hist_wx$DATE, format = "%Y-%m-%d")
```

```
  return(hist_wx)
```

```
}
```

---

 get\_gsod\_stn

---

*Obtain Global Summary of Day data*


---

**Description**

This function will grab available GSOD data for stations and years for which the data is available.

**Usage**

```
get_gsod_stn()
```

**Author(s)**

Daniel R. Fuka

**Examples**

```
# Example of downloading data for ADDIS ABABA
```

```
## Not run: get_gsod_stn()
```

```
# Please enter a temp directory ....
```

```
addis
```

```
# What is the station name you want to look for?
```

```

ADDIS\ A
#      USAF  WBAN      STATION.NAME ....
#13582 634500 99999 ADDIS ABABA-BOLE ....
#Good Enough?
y
#Given the start and end years, ....
n
#Date range available is 1957 to 2011 , what start year would you like?
1990
#Date range available is 1957 to 2011 , what end year would you like?
1992
#Good enough?
y

## End(Not run)

## The function is currently defined as
function(){
  tmpdir=readline("Please enter a temp directory to store the datafiles in, remember this directory
when parsing with build_gsod_forcing_data()? \n")
  dir.create(tmpdir)
  stnlist=GSOD_history
  ANSWER="n"
  while(ANSWER=="n"){
    stnreq=readline("What is the station name you want to look for? \n")
    test=grep(stnreq,stnlist$STATION.NAME)
    print(stnlist[test,])
    ANSWER <- readline("Good enough? ")
  }
  ANSWER <- readline("Given the start and end years, would you like to
update the stnlist file from NDCD? (may take a few minutes) ")
  ## a better version would check the answer less cursorily, and
  ## perhaps re-prompt
  if (substr(ANSWER, 1, 1) == "y")
    stnlist=read.csv(url("
ftp://ftp.ncdc.noaa.gov/pub/data/gsod/ish-history.csv"))
#
# Add one more request to make sure the number of years is ok as some
# stations have really long histories.
#
#
  working_tmp <- stnlist$STATION.NAME
  has_stn<- working_tmp %in% grep(stnreq, working_tmp, value=TRUE)
  dl_wbans=subset(stnlist, has_stn , USAF:END)
  start_year=min(trunc(subset(stnlist, has_stn, BEGIN)/10000))
  end_year=max(trunc(subset(stnlist, has_stn , END)/10000))
  start_year_rec<- readline(paste("Date range available
is",start_year,"to",end_year,", what start year would you like?"))
  end_year_rec<- readline(paste("Date range available
is",start_year,"to",end_year,", what end year would you like?"))
  for (wbans in row.names(dl_wbans)){
    start_year=max(start_year_rec,substr(dl_wbans[wbans,"BEGIN"],1,4))
    end_year=min(end_year_rec,substr(dl_wbans[wbans,"END"],1,4))
  }
}

```



```

    print(c(start_year,end_year))
    if(start_year > end_year) next()
    for (year in start_year:end_year){
      dlurl=paste("ftp://ftp.ncdc.noaa.gov/pub/data/gsod/
",year,"/",dl_wbans[wbans,1],"-",dl_wbans[wbans,2],"-",year,".op.gz"
,sep="")

      dlfile=paste(tmpdir,"/",dl_wbans[wbans,1],"-",dl_wbans[wbans,2],"-
",year,".op.gz",sep="")
      print(dlurl)
      try(download.file(dlurl,dlfile),silent=T)

    }
  }
  files=dir(tmpdir,"*op.gz")
  print(files)
  ANSWER <- readline("Good enough? ")
}

```

---

get\_usgs\_gage

*A function to grab daily stream flow measurements from the USGS waterdata.usgs.gov server.*

---

## Description

A function to grab daily stream flow measurements from the USGS waterdata.usgs.gov server.

## Usage

```
get_usgs_gage(flowgage_id,begin_date="1979-01-01",end_date="2013-01-01")
```

## Arguments

flowgage_id	flowgage_id Gage is a TEXT/String rather than numeric or the query will fail on gages with leading 0s
begin_date	begin_date begin_date is a TEXT/String for the start date for the data you want. Default is the beginning of the cfsr dataset.
end_date	end_date end_date is a TEXT/String for the end date for the data you want. Default is in the future sometime

## Value

area	The area above the gage [km2]
declat	Decimal Lat
declon	Decimal Lon
flowdata	Dataframe with the historical flow data - flow is in cubic meters per day

**Author(s)**

Daniel R Fuka

**Examples**

```
flowgage_id="04216500"  
flowgage=get_usgs_gage(flowgage_id)  
  
## The function is currently defined as
```

---

GroundHeat

*Heat conducted to the bottom of a snowpack*

---

**Description**

The heat conducted to the bottom of a snowpack, assumed constant [kJ m<sup>-2</sup> d<sup>-1</sup>]

**Usage**

GroundHeat()

**Author(s)**

M. Todd Walter

**Examples**

```
##---- Should be DIRECTLY executable !! ----  
##-- ==> Define data, use random,  
##--or do help(data=index) for the standard data sets.  
  
## The function is currently defined as  
function(){  
# the heat conducted to the bottom of a snowpack, assumed constant [kJ m-2 d-1]  
  
return(173)  
}
```

---

GSOD_history	<i>ish-history.txt</i>
--------------	------------------------

---

**Description**

A station list to be used with the GSOD data files, showing the names and locations for each station.  
Note: Global summary of day contains a subset of the stations listed in this station history.

**Usage**

```
data(GSOD_history)
```

**Format**

A data frame with 30558 observations on the following 12 variables.

USAF a numeric vector

WBAN a numeric vector

STATION.NAME a factor with levels

CTRY a factor with levels

FIPS a factor with levels

STATE a factor with levels

CALL a factor with levels

LAT a numeric vector

LON a numeric vector

ELEV..1M. a numeric vector

BEGIN a numeric vector

END a numeric vector

**Source**

<ftp://ftp.ncdc.noaa.gov/pub/data/gsod/ish-history.csv>

**Examples**

```
data(GSOD_history)
```

---

hydrograph

*Plot a hydrograph*


---

### Description

Creates a hydrograph with one or two streamflow data sets, and can include a precipitation hydrograph.

### Usage

```
hydrograph(input = matrix(ncol = 2, nrow = 2), streamflow = input[, 2],
  timeSeries = input[, 1], streamflow2 = NULL, precip = NULL, begin = 1,
  endindex = length(streamflow), P.units = "", S.units = "",
  S1.col = "black", S2.col = "red", stream.label = "Streamflow",
  streamflow3 = NULL, streamflow4 = NULL, precip2 = NULL)
```

### Arguments

input	This is a data frame with at least two columns of data First column: Must be a time/date series Second column: If including precip, precip. Otherwise, streamflow - AND do not include any other columns Third column: (Only include if precip is in the second column) First streamflow dataset Forth column: (optional) Second streamflow dataset
streamflow	vector of streamflow values - not necessary if using "input"
timeSeries	vector of time or date values - not necessary if using "input"
streamflow2	optional vector of streamflow values - not necessary if using "input"
precip	vector of precipitation values - not necessary if using "input"
begin	If you don't want to graph the whole data set, include the index where you would like to start
endindex	If you don't want to graph the whole dataset, include the index where you would like to end
P.units	Precipitation units (character)
S.units	Streamflow units (character). Users who have volumetric flow will have the superscript correctly formatted if they enter "m3s" or "ft3s"
S1.col	color of the line representing streamflow
S2.col	color of the line representing second streamflow data
stream.label	character string - Label for vertical axis
streamflow3	optional vector of additional streamflow values
streamflow4	optional vector of additional streamflow values
precip2	optional vector of a second precip gage data if you are interested in comparing precip inputs

**Warning**

The date series should be continuous and evenly spaced. If not, the dates will not line up accurately on the x-axis.

**Note**

This function can now take NA values. If you chose to use the input argument but are not including precipitation, input must only have two columns - you will need to add the second streamflow dataset using the "streamflow2" argument.

**Author(s)**

Josephine Archibald

**Examples**

```
data(OwascoInlet)
head(OwascoInlet)
hydrograph(OwascoInlet)
```

---

Longwave

*Daily Longwave Radiation*

---

**Description**

Daily longwave radiation based on the Stefan-Boltzman equation [kJ m<sup>-2</sup> d<sup>-1</sup>]

**Usage**

```
Longwave(emissivity, temp)
```

**Arguments**

emissivity	emissivity [-]
temp	temperature of the emitting body [C]

**Author(s)**

M. Todd Walter

---

Lumped\_VSA\_model      *Lumped Variable Source Area (VSA) Watershed Model*

---

### Description

This model calculates streamflow and approximate saturated area percentage contributing to overland flow. It was developed in saturation-excess dominated watersheds, and is based on the Thornthwaite-Mather water budget and SCS Curve Number approach for overland runoff.

### Usage

```
Lumped_VSA_model(dateSeries, P, Tmax, Tmin, Depth = NULL, SATper = NULL, AWCper = NULL,
percentImpervious = 0, no_wet_class = 10, Tp = 5, latitudeDegrees = 42.38, albedo = 0.23,
StartCond = "avg", PETin = NULL, AWC = Depth * AWCper, SAT = Depth * SATper, SW1 = NULL,
BF1 = 1, PETcap = 5, rec_coef = 0.1, Se_min = 78, C1 = 3.1, Ia_coef = 0.05,
PreviousOutput = NULL, runoff_breakdown = RunoffBreakdown(Tp, HrPrcDelay = (Tp/2 - 4)))
```

### Arguments

dateSeries	Daily date series in the format "2013-05-21"
P	Rain + Snowmelt (mm)
Tmax	Maximum daily T (C)
Tmin	Minimum daily T (C)
Depth	Average watershed soil depth (mm) Not needed if SAT and AWC depth entered directly
SATper	Porosity of the soil (volumetric fraction, 0-1) Not needed if SAT (porosity depth) entered directly
AWCper	Available water capacity, Field capacity - wilting point (volumetric fraction, 0-1) Not needed if AWC entered directly
percentImpervious	Percent of the watershed that is impervious (percentage, 0-100)
no_wet_class	Number of wetness classes to distribute runoff over. Default is 10.
Tp	Time to peak of hydrograph (hours)
latitudeDegrees	latitude (degrees)
albedo	Average surface albedo, defaults to average 0.23
StartCond	Watershed conditions before first day of run (options are "wet", "dry", "avg")
PETin	# User has the option to enter PET values (mm/day), otherwise this will be estimated from Priestley-Taylor equation, estimating radiation from temperature
AWC	# AWC depth (mm)
SAT	Porosity depth (mm)
SW1	Soil water on the first day (depth, mm)

BF1	Baseflow on the first day (mm/day)
PETcap	Cutoff for maximal PET allowed per day (mm)
rec_coef	Baseflow recession coefficient
Se_min	Minimal daily CN S value. (mm)
C1	Coefficient relating daily Curve Number S to soil water
Ia_coef	Initial abstraction coefficient for CN-equation. (range ~ 0.05 - 0.2)
PreviousOutput	If the model is run repeatedly, previous output can be provided so that the model only needs to calculate from that point forward.
runoff_breakdown	The proportion of runoff that reaches the outlet on a given day after the storm event. Can be calculated from Tp

### Details

The model expects continuous input on a daily time-step, since the soil-water is calculated over time, and affects the amount of runoff that will be generated after a storm. Also, note that precipitation values are actually Rain + Snowmelt (mm). Users can use the snowmelt function to determine this if needed.

### Value

Returns a data frame with modeled streamflow, baseflow, ET, and maximum wetness class generating runoff for all dates. Soil water and other modeled intermediate results are also returned. All flow values (modeled\_flow, baseflow, OverlandFlow, ShallowInterflow, totQ, quickflow\_combined, impervRunoff, excess) are in depth of flow per day (mm/d)

### Warning

This function cannot handle NA values in input, and can only be run for a daily time-step. If  $T_x < T_n$  for any day, this will produce an error. Currently, the crop coefficients used to estimate PET are specific for deciduous northeastern USA.

### Note

This function is under development and might change substantially in further versions.

### Author(s)

Josephine Archibald

### References

Archibald, J.A., B.P. Buchanan, D.R. Fuka, C.B. Georgakakos, S.W. Lyon, M.T. Walter. A simple, regionally parameterized model for predicting nonpoint source areas in the Northeastern US. Submitted to Journal of Hydrology: Regional Studies

Schneiderman EM, Steenhuis TS, Thongs DJ, Easton ZM, Zion MS, Neal AL, Mendoza GF, Walter MT. 2007. Incorporating variable source area hydrology into a curve-number-based watershed model. Hydrological Processes. 21: 3420-3430

Shaw, SB, MT Walter. 2009. Improving runoff risk estimates: Formulating runoff as a bivariate process using the SCS curve number method. *Water Resources Research*. 45

Thornthwaite CW, JR Mather. 1957 Instructions for computing potential evapotranspiration and water balance. *Publ Climatol* 3: 183-311

United States Department of Agriculture (USDA). (1986). SCS publication Technical Release 55: Urban Hydrology for Small Watersheds. Natural Resources Conservation Service

Weiler K. Unpublished. Determination of the Linear Bedrock Coefficient From Historical Flow Data

### See Also

PET\_fromTemp

### Examples

```
data(OwascoInlet)
# First get rain and snow-melt input:
rsm <- SnowMelt(Date=OwascoInlet$date, precip_mm=OwascoInlet$P_mm, Tmax_C=OwascoInlet$Tmax_C,
Tmin_C=OwascoInlet$Tmin_C, lat_deg=42.66)
# Calculate streamflow based on watershed characteristics:
Results <- Lumped_VSA_model(dateSeries = OwascoInlet$date, P = rsm$Rain_mm+rsm$SnowMelt_mm,
Tmax=OwascoInlet$Tmax_C, Tmin = OwascoInlet$Tmin_C, latitudeDegrees=42.66, Tp = 5.8, Depth = 2010,
SATper = 0.27, AWCper = 0.13, StartCond = "wet")
# View results:
hydrograph(streamflow=ConvertFlowUnits(cms=OwascoInlet$Streamflow_m3s, WA=106, AREAunits="mi2"),
timeSeries=OwascoInlet$date, streamflow2=Results$modeled_flow, precip=rsm$Rain_mm+rsm$SnowMelt_mm)
```

---

NetRad

*Daily Net Radiation*

---

### Description

Daily net radiation [kJ m<sup>-2</sup> d<sup>-1</sup>]

### Usage

```
NetRad(lat, Jday, Tx, Tn, albedo = 0.18, forest = 0, slope = 0,
aspect = 0, airtemp = (Tn+Tx)/2, cloudiness = "Estimate",
surfemissivity = 0.97, surftemp = (Tn+Tx)/2, units = "kJm2d",
AEparams=list(vp=NULL, opt="linear"))
```

### Arguments

lat	latitude. Default is radians, but will automatically convert from degrees if value is larger than pi/2 or less than -pi/2
Jday	Day of the year [-]
Tx	Max temperature [C]



Tn	Min temperature [C]
albedo	Albedo (0-1) [-]
forest	Forest canopy or shade cover (0-1) [-]
slope	Slope [radians]
aspect	Ground aspect [radians from north]
airtemp	Average air temp [C]
cloudiness	Cloudiness percentage (0-1) [-] Default value will estimate cloudiness based on daily temperature difference.
surfemissivity	Surface emissivity [-]
surftemp	Surface temperature [C]
units	Units of the result. Defaults to "kJm2d" which returns result in kJ/m2/d, the other option is "Wm2", which will make output in W/m2
AEparams	Atmospheric Emissivity options. Defaults to linear approximation. To use Brutsaert option, include vapor pressure values (kPa)

**Author(s)**

M. Todd Walter, Josephine Archibald

**Examples**

```
NetRad(42.4*pi/180, Jday=55, Tx=2, Tn=(-3))
```

---

OwascoInlet

*Owasco Inlet data*

---

**Description**

Streamflow and temperature data for Owasco Inlet from USGS and NRCC

**Usage**

```
data(OwascoInlet)
```

**Format**

A data frame with 888 observations on the following 6 variables.

date a Date

P\_mm a numeric vector

Streamflow\_m3s a numeric vector

baseflow\_m3s a numeric vector

Tmax\_C a numeric vector

Tmin\_C a numeric vector

**Source**

<http://www.usgs.gov/> <http://www.nrcc.cornell.edu/>

**References**

Data taken from USGS ([www.usgs.gov](http://www.usgs.gov/)) and NRCC ([http://www.nrcc.cornell.edu](http://www.nrcc.cornell.edu/))

**Examples**

```
data(OwascoInlet)
## maybe str(OwascoInlet) ; plot(OwascoInlet) ...
```

---

PET\_fromTemp

*Priestley-Taylor Potential Evapotranspiration from temperature*

---

**Description**

Calculates potential Evapotranspiration (in meters) based on the Priestley-Taylor equation (1972). We use an estimation of net radiation based on temperature data.

**Usage**

```
PET_fromTemp(Jday, Tmax_C, Tmin_C, lat_radians,
AvgT = (Tmax_C + Tmin_C)/2, albedo = 0.18, TerrestEmiss = 0.97,
aspect = 0, slope = 0, forest = 0, PTconstant=1.26,
AParams=list(vp=NULL, opt="linear"))
```

**Arguments**

Jday	Day of the year
Tmax_C	Maximum daily temperature (degrees C)
Tmin_C	Minimum daily temperature (degrees C)
lat_radians	latitude (radians = decimal degrees*pi/180)
AvgT	Average daily temperature (degrees C) (if not known, will be taken as the averages of the daily extremes)
albedo	(-) average surface albedo. Can be expressed as a single value, or as a vector with the same length as Jday, Tmax_C and Tmin_C
TerrestEmiss	(-) Surface Emissivity - defaults to 0.97
aspect	(radians) Surface aspect
slope	(radians) average slope
forest	(-) Forest or shade cover (0-1). This modifies the amount of solar radiation reaching the location of interest. It should always set to zero for landscape-wide processes regardless of the amount of forest present. Only change this if calculating PET under a canopy.
PTconstant	(-) Priestley-Taylor Constant, often 1.26
AParams	Atmospheric Emissivity options. Defaults to linear approximation. To use Brutsaert option, include vapor pressure values (kPa)

**Value**

PET (potential evapotranspiration) in m

**Note**

We are assuming that the Ground heat flux on a daily time-step is zero

**Author(s)**

Josephine Archibald, M. Todd Walter

**References**

Archibald, J.A. and M. T. Walter, 2013. Do energy-based PET models require more input data than T-based models? - An evaluation at four humid FluxNet sites. *Journal of the American Water Resources Association (JAWRA)*

Brutsaert, W., 1975. On a Derivable Formula for Long-Wave Radiation from Clear Skies. *Water Resources Research* 11(5):742-744.

Priestley and Taylor, 1972. On the assessment of surface heat flux and evaporation using large-scale parameters. *Mon. Weath. Rev.* 100: 81-92

**Examples**

```
data(OwascoInlet)
head(OwascoInlet)
attach(OwascoInlet)
PETapprox <- PET_fromTemp(Jday=(1+as.POSIXlt(date)$yday), Tmax_C=Tmax_C,
  Tmin_C=Tmin_C, lat_radians=42.45*pi/180)
plot(PETapprox*1000~date, type="l")
detach(OwascoInlet)
```

---

PotentialSolar

*Potential Solar Radiation*

---

**Description**

Potential solar radiation at the edge of the atmosphere [kJ m<sup>-2</sup> d<sup>-1</sup>]

#lat: latitude [rad] #Jday: Julian date or day of the year [day]

**Usage**

```
PotentialSolar(lat, Jday)
```

**Arguments**

lat                    latitude in radians

Jday                  Day of the year

**Author(s)**

M. Todd Walter

PotSolarInst

*Instantaneous Potential Solar Radiation***Description**

Potential Solar Radiation at a particular time of day. Defaults to W/m<sup>2</sup>, can also report in kJ/m<sup>2</sup>/d if units set to kJ/m<sup>2</sup>/d

**Usage**

```
PotSolarInst(Jday, hour = 12, lat = 42.44 * pi/180, sunrise = NULL,
  sunset = NULL, SolarNoon = mean(c(sunrise, sunset)), units = "Wm2",
  latUnits = "unknown")
```

**Arguments**

Jday	Day of the year [-]
hour	Time of the day in hours [0-24 hr]
lat	latitude. Default is radians, but will automatically convert from degrees if value is larger than 1.62 or less than -1.62
sunrise	Time of sunrise used to calculate solar noon [0-24 hr]
sunset	Time of sunset used to calculate solar noon [0-24 hr]
SolarNoon	Time of solar noon. Can be calculated from sunrise and sunset times. [hr]
units	Units of the result. Defaults to W/m <sup>2</sup>
latUnits	Latitude units can be explicitly stated here, options are 'radians', 'degrees' or default is 'unknown', which will assume radians unless the absolute value of lat is greater than pi/2

**Author(s)**

Josephine Archibald

**References**

Lawrence Dingman. Physical Hydrology. Waveland Press, Inc. Illinois, 2002.

**Examples**

```
PotSolarInst(Jday=150, hour = 15, lat = 42.44, SolarNoon = 12.5)
PotSolarInst(Jday=c(1,50,100,150), hour = c(9,10,12,17), lat = -pi/4, SolarNoon = 12.5)
```

---

PTpet                      *Priestley-Taylor potential evapotranspiration*

---

**Description**

Returns potential evapotranspiration in m/day caculated with the Priestley-Taylor equation

**Usage**

PTpet(Rn, T\_C, PTconstant = 1.26)

**Arguments**

Rn	Net daily radiation [kJ/m2/d]
T_C	Average daily temperature [C]
PTconstant	Priestley-Taylor constant, usually 1.26 [-]

**Author(s)**

Josephine Archibald

**References**

Priestley and Taylor (1972). On the assessment of surface heat flux and evaporation using large-scale parameters. Mon. Weath. Rev. 100: 81-92

---

RainHeat                      *Heat from Rain*

---

**Description**

Temperature added to the land from heat exchange with rain (usually in the context of snowmelt) [kJ m-2 d-1]

**Usage**

RainHeat(airtemp, rain)

**Arguments**

airtemp	average dailiy air temperature [C]
rain	depth of rainfall [m]

**Examples**

RainHeat(20,0.01)

---

RunoffBreakdown      *Daily overland runoff breakdown*

---

**Description**

Determines the percentage of runoff that occurs on each day following a storm event, based on time to peak of a watershed.

**Usage**

```
RunoffBreakdown(Tp_hr, a = 4.5, HrPrcDelay = 4, numDaysReturn = 5)
```

**Arguments**

Tp_hr	Time to peak (hr).
a	The ratio of the time to recession to the time to peak. Default is 4.5
HrPrcDelay	Lag time (hr)
numDaysReturn	Number of days to return

**Value**

returns a vector corresponding to the amount of runoff per day

**Author(s)**

J Archibald

**Examples**

```
RunoffBreakdown(5)
```

---

SatVaporDensity      *Saturated Vapor Density*

---

**Description**

Saturated vapor density [kg/m<sup>3</sup>] at a given temperature

**Usage**

```
SatVaporDensity(T_C)
```

**Arguments**

T_C	temperature [C]
-----	-----------------

**Author(s)**

Josephine Archibald, M. Todd Walter

**References**

Campbell, Gaylon S. An Introduction to Environmental Biophysics. New York: Springer-Verlag, 1977.

**Examples**

```
SatVaporDensity(seq(-5,40, by=5))
```

---

SatVaporPressure	<i>Saturated Vapor Pressure</i>
------------------	---------------------------------

---

**Description**

Calculates the saturated vapor pressure (kPa) at a particular temperature (degrees Celcius)

**Usage**

```
SatVaporPressure(T_C)
```

**Arguments**

T\_C

**Author(s)**

Josephine Archibald

**References**

Tetens (1930) : Tetens, V.O (1930) Uber einige meteorologische. Begriffe, Zeitschrift fur Geophysik. 6, 297-309

---

SatVapPresSlope	<i>Slope of the relationship between Saturation Vapor Pressure and Temperature</i>
-----------------	--

---

**Description**

Calculates the slope of the Saturation Vapor Pressure vs T (kPa/K)

**Usage**

```
SatVapPresSlope(temp_C)
```

**Arguments**

temp_C	Air temperature in degrees C
--------	------------------------------

**Author(s)**

Josephine Archibald

**References**

Dingman, S. Lawrence. 2002. Physical Hydrology. Waveland Press Inc.

**Examples**

```
## The function is currently defined as
function (temp_C)
{
  (2508.3/(temp_C + 237.3)^2) * exp(17.3 * temp_C/(temp_C +
    237.3))
}
```

---

SensibleHeat	<i>Sensible Heat Exchange</i>
--------------	-------------------------------

---

**Description**

Sensible heat exchange between a surface and the surrounding air [kJ m<sup>-2</sup> d<sup>-1</sup>]

**Usage**

```
SensibleHeat(surftemp, airtemp, wind)
```



**Arguments**

surftemp	surface temperature [C]
airtemp	average dailiy air temperature [C]
wind	average daily windspeed [m/s]

**Author(s)**

M. Todd Walter

**Examples**

```
##---- Should be DIRECTLY executable !! ----
##-- ==> Define data, use random,
##--or do help(data=index) for the standard data sets.

## The function is currently defined as
function(surftemp,airtemp,wind){
# sensible heat exchange between a surface and the surrounding air [kJ m-2 d-1]

#surftemp: surface temperature [C]
#airtemp: average dailiy air temperature [C]
#wind: average daily windspeed [m/s]

latentht<-2500 #latent heat of vaporization [kJ kg-1]

heatcapacity<-1.25 #approx. heat capacity of air [kJ m-3 C-1]

windfunction<-5.3*(1+wind)

return(86400*heatcapacity*(surftemp-airtemp)*windfunction/latentht)
}
```

---

setup\_swatcal

*A function to setup a swat calibration*

---

**Description**

A function to setup a swat calibration, building the template \*.unixorig template files, and cleaning up oddities we find in the default input files. (eg file.cio has cst.cst file which appears to be a bug)

**Usage**

```
setup_swatcal(change_params)
```

**Arguments**

change\_params

**Author(s)**

Daniel Fuka drf28@cornell.edu

**Examples**

```
##---- Should be DIRECTLY executable !! ----
##-- ==> Define data, use random,
##--or do help(data=index) for the standard data sets.

## The function is currently defined as
function(change_params){
  library(operators)
  for(ft in unique(change_params$filetype)){
    print(ft)
    files=list.files(,paste(ft,"$",sep=""))
    for (file in files) {
      junk%<%file
      junk=gsub("\r", "", junk)
      file_swatcal=paste(file, ".unixorig", sep="");
      cat(junk, file=file_swatcal, sep="\n")
    }
  }
}
```

---

slopefactor

*Slope Factor*

---

**Description**

Adjusts solar radiation for land slope and aspect relative to the sun, 0=level ground

**Usage**

```
slopefactor(lat, Jday, slope, aspect)
```

**Arguments**

lat	latitdue [rad]
Jday	Julian date or day of the year [day]
slope	slope of the ground [rad]
aspect	ground aspect [rad from north]

**Author(s)**

M. Todd Walter, J. Archibald

**References**

Monteith and Unsworth (1990) Principles of Environmental Physics, 2nd ed. Chapter 5

SnowMelt

*Snow Melt Calculator***Description**

Calculates snow pack accumulation and melt

**Usage**

```
SnowMelt(Date, precip_mm, Tmax_C, Tmin_C, lat_deg, slope = 0,
  aspect = 0, tempHt = 1, windHt = 2, groundAlbedo = 0.25,
  SurfEmissiv = 0.95, windSp = 2, forest = 0, startingSnowDepth_m = 0,
  startingSnowDensity_kg_m3=450)
```

**Arguments**

Date	Vector of dates (class Date or character) in this format: Y-m-d
precip_mm	Precipitation in mm
Tmax_C	Daily maximum temperature (degrees C)
Tmin_C	Daily minimum temperature (degrees C)
lat_deg	Degrees latitude
slope	Overall slope of area of interest
aspect	Aspect of the area of interest
tempHt	height of temperature measurements (m)
windHt	height of wind measurements (m)
groundAlbedo	Ground Albedo, 0-1 (-)
SurfEmissiv	Surface Emissivity, 0-1 (-)
windSp	Wind speed - either a vector of measured values or a single value of average wind speed for the site (m/s)
forest	Forest cover (shade) - use this only when determining snowmelt under a canopy, 0-1 (-)
startingSnowDepth_m	The depth of the snow pack initially (m)
startingSnowDensity_kg_m3	The density of snow on the ground on the first day (kg/m3)

**Value**

This will return a 10 column data frame with nrow = length of input weather data. Column names are : Date, MaxT\_C, MinT\_C, Precip\_mm, Rain\_mm, SnowfallWatEq\_mm, SnowMelt\_mm, NewSnow\_m, SnowDepth\_m, SnowWaterEq\_mm

**Warning**

This function cannot handle NA values in input, and can only be run for continuous daily data. For data-sets missing values, run discontinuous segments separately.

**Author(s)**

Josephine Archibald, M. Todd Walter

**References**

Walter, M.T. , E.S. Brooks, D.K. McCool, L.G. King, M. Molnau, J. Boll. 2005. Process-based snowmelt modeling: Does it require more input data than temperature-index modeling? *Journal of Hydrology* 300(1-4): 65-75

**Examples**

```
##
data(OwascoInlet)
sm <- SnowMelt(Date=OwascoInlet$date, precip_mm=OwascoInlet$P_mm,
  Tmax_C=OwascoInlet$Tmax_C, Tmin_C=OwascoInlet$Tmin_C, lat_deg=42)
summary(sm)
```

---

 SoilStorage

---

*Calculating S in the Curve Number Equation from soil water content*


---

**Description**

This function calculates S, used in the SCS-CN equation, from the water content of the soil.

**Usage**

```
SoilStorage(S_avg, field_capacity, soil_water_content, porosity)
```

**Arguments**

S_avg	Average S, as used normally in the CN equation, calculated from the curve number, which is based on land-use. This is in units of depth, often mm or inches
field_capacity	field capacity - the amount of water that a soil can hold after drainage. (fraction)
soil_water_content	Soil water content (fraction) on a given day.
porosity	Saturated water content, approximately equal to the porosity of a soil (fraction)

**Note**

This equation is not the same as the one used in SWAT. It was given to Dr. Todd Walter by Dr. Keith E. Saxton

**Author(s)**

Josephine Archibald

**See Also**

SoilStorageSWAT

**Examples**

```
SoilStorage(S_avg=120, field_capacity=0.2, soil_water_content=0.1, porosity=0.3)
```

Solar

*Solar Radiation***Description**Solar radiation at the ground surface [kJ m<sup>-2</sup> d<sup>-1</sup>]**Usage**

```
Solar(lat, Jday, Tx, Tn, albedo=0.2, forest=0, slope=0, aspect = 0,
      units="kJm2d", latUnits = "unknown", printWarn=TRUE)
```

**Arguments**

lat	latitude [rad]
Jday	Julian date or day of the year [day]
Tx	maximum daily temperature [C]
Tn	minimum daily temperature [C]
albedo	surface albedo or reflectivity [-]
forest	forest or vegetation cover [-]
slope	slope of the ground [rad]
aspect	ground aspect [rad from north]
units	Units of the result. Defaults to kJ/m <sup>2</sup> /d, changing this to "Wm <sup>2</sup> " will make output in W/m <sup>2</sup>
latUnits	Latitude units can be explicitly stated here, options are 'radians', 'degrees' or default is 'unknown', which will assume radians unless the absolute value of lat is greater than pi/2
printWarn	Will print a warning about latitude if set to TRUE

---

solarangle

*Solar Angle*


---

**Description**

Angle of solar inclination from horizontal at solar noon [rad]

**Usage**

```
solarangle(lat, Jday)
```

**Arguments**

```
lat          latitdue [rad]
Jday         Julian date or day of the year [day]
```

**Examples**

```
##---- Should be DIRECTLY executable !! ----
##-- ==> Define data, use random,
##--or do help(data=index) for the standard data sets.

## The function is currently defined as
function(lat,Jday){
# angle of solar inclination from horizontal at solar noon [rad]

#lat: latitdue [rad]
#Jday: Julian date or day of the year [day]

# solar declination [rad]
dec<-declination(Jday)

return(asin(sin(lat)*sin(dec)+cos(lat)*cos(dec)*cos(0)))
}
```

---

SWAT2005

*This runs the SWAT2005 executable in the current directory.*


---

**Description**

This function runs the SWAT2005 executable in the current directory.

**Usage**

```
SWAT2005(hist_wx,elev,rch)
```

**Arguments**

hist_wx	Describe hist_wx
elev	Describe elev
rch	Describe rch

**Author(s)**

Daniel R. Fuka

---

swat\_general

*All files required for a general SWAT run.*


---

**Description**

All files required for a general SWAT run. This initialization has 3 subbasin init, with 3 HRUs per subbasin.

**Usage**

data(swat\_general)

**Format**

The format is: List of 74 \$ 000010000.pnd: chr [1:47] ".pnd file Subbasin: 2 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT interface" "Pond inputs:" " 0 | PND\_FR : Fraction of sub-basin area that drains into ponds. The value for PND\_FR should be between 0.0 and"| \_\_truncated\_\_ " 0 | PND\_PSA: Surface area of ponds when filled to principal spillway [ha]" ... \$ 000010000.rte: chr [1:23] ".rte file Subbasin: 2 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT interface" "9.3638387596228 | CH\_W2 : Main channel width [m]" ".304414007785773 | CH\_D : Main channel depth [m]" "0.03293296544532192 | CH\_S2 : Main channel slope [m/m]" ... \$ 000010000.sub: chr [1:64] ".sub file Subbasin: 1 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT interface" "15 | SUB\_KM : Subbasin area [km2]" "" "Climate in subbasin" ... \$ 000010000.swq: chr [1:30] ".swq file Subbasin: 2 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT interface" "Nutrient (QUAL2E parameters)" " 1 | RS1: Local algal settling rate in the reach at 20 degC [m/day]" " 0.05 | RS2: Benthic (sediment) source rate for dissolved phosphorus in the reach at 20 degC [mg dissolved P/[m"| \_\_truncated\_\_ ... \$ 000010000.wgn: chr [1:19] ".wgn file Subbasin: 2 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT interface" " LATITUDE = 20.00 LONGITUDE = 0.00" " ELEV [m] =2000.00" " RAIN\_YRS = 10.00" ... \$ 000010000.wus: chr [1:11] ".wus file Subbasin: 2 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT interface" "" "" " 0.0 0.0 0.0 0.0 0.0 0.0" ... \$ 000010001.chm: chr [1:21] ".chm file Subbasin: 2 HRU: 1 Luse: SHRB Soil: K115-2ab-5212 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inte"| \_\_truncated\_\_ "Soil Nutrient Data" " Soil Layer : 1 2 3 4 5 6 7 8 "| \_\_truncated\_\_ " Soil NO3 [mg/kg] : 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.00 "| \_\_truncated\_\_ ... \$ 000010001.gw : chr [1:14] ".gw file Subbasin: 2 HRU: 1 Luse: SHRB Soil: K115-2ab-5212 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inter"| \_\_truncated\_\_ " 0.5 | SHALLST : Initial depth of water in the shallow aquifer [mm]" " 1000 |

DEEPST : Initial depth of water in the deep aquifer [mm]" " 31 | GW\_DELAY : Groundwater delay [days]" ... \$ 000010001.hru: chr [1:25]" .hru file Subbasin: 2 HRU: 1 Luse: SHRB Soil: K115-2ab-5212 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inte"| \_\_truncated\_\_ "0.347771714415945 | HRU\_FR : Fraction of subbasin area contained in HRU" " 90 | SLSUBBSN : Average slope length [m]" "0.0260677934475378 | HRU\_SLP : Average slope stepness [m/m]" ... \$ 000010001.mgt: chr [1:33]" .mgt file Subbasin: 2 HRU: 1 Luse: SHRB Soil: K115-2ab-5212 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inte"| \_\_truncated\_\_ " 0 | NMGT: Management code" "Initial Plant Growth Parameters" " 0 | IGRO: Land cover status: 0-none growing; 1-growing" ... \$ 000010001.sol: chr [1:19]" .sol file Subbasin: 2 HRU: 1 Luse: SHRB Soil: K115-2ab-5212 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inte"| \_\_truncated\_\_ "Soil Name: K115-2ab-5212" "Soil Hydrologic Group: C" "Maximum rooting depth(m): 580.00" ... \$ 000010002.chm: chr [1:21]" .chm file Subbasin: 2 HRU: 2 Luse: SHRB Soil: K130-2bc-4832 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inte"| \_\_truncated\_\_ "Soil Nutrient Data" " Soil Layer : 1 2 3 4 5 6 7 8 "| \_\_truncated\_\_ " Soil NO3 [mg/kg] : 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.00 "| \_\_truncated\_\_ ... \$ 000010002.gw : chr [1:14]" .gw file Subbasin: 2 HRU: 2 Luse: SHRB Soil: K130-2bc-4832 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inter"| \_\_truncated\_\_ " 0.5 | SHALLST : Initial depth of water in the shallow aquifer [mm]" " 1000 | DEEPST : Initial depth of water in the deep aquifer [mm]" " 31 | GW\_DELAY : Groundwater delay [days]" ... \$ 000010002.hru: chr [1:25]" .hru file Subbasin: 2 HRU: 2 Luse: SHRB Soil: K130-2bc-4832 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inte"| \_\_truncated\_\_ "0.351068667576475 | HRU\_FR : Fraction of subbasin area contained in HRU" " 90 | SLSUBBSN : Average slope length [m]" "0.0246108923185198 | HRU\_SLP : Average slope stepness [m/m]" ... \$ 000010002.mgt: chr [1:33]" .mgt file Subbasin: 2 HRU: 2 Luse: SHRB Soil: K130-2bc-4832 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inte"| \_\_truncated\_\_ " 0 | NMGT: Management code" "Initial Plant Growth Parameters" " 0 | IGRO: Land cover status: 0-none growing; 1-growing" ... \$ 000010002.sol: chr [1:19]" .sol file Subbasin: 2 HRU: 2 Luse: SHRB Soil: K130-2bc-4832 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inte"| \_\_truncated\_\_ "Soil Name: K130-2bc-4832" "Soil Hydrologic Group: C" "Maximum rooting depth(m): 460.00" ... \$ 000010003.chm: chr [1:21]" .chm file Subbasin: 2 HRU: 3 Luse: GRAS Soil: Vp38-3a-5345 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inter"| \_\_truncated\_\_ "Soil Nutrient Data" " Soil Layer : 1 2 3 4 5 6 7 8 "| \_\_truncated\_\_ " Soil NO3 [mg/kg] : 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.00 "| \_\_truncated\_\_ ... \$ 000010003.gw : chr [1:14]" .gw file Subbasin: 2 HRU: 3 Luse: GRAS Soil: Vp38-3a-5345 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT interf"| \_\_truncated\_\_ " 0.5 | SHALLST : Initial depth of water in the shallow aquifer [mm]" " 1000 | DEEPST : Initial depth of water in the deep aquifer [mm]" " 31 | GW\_DELAY : Groundwater delay [days]" ... \$ 000010003.hru: chr [1:25]" .hru file Subbasin: 2 HRU: 3 Luse: GRAS Soil: Vp38-3a-5345 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inter"| \_\_truncated\_\_ "0.301159618008454 | HRU\_FR : Fraction of subbasin area contained in HRU" " 90 | SLSUBBSN : Average slope length [m]" "0.0277888825969753 | HRU\_SLP : Average slope stepness [m/m]" ... \$ 000010003.mgt: chr [1:33]" .mgt file Subbasin: 2 HRU: 3 Luse: GRAS Soil: Vp38-3a-5345 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inter"| \_\_truncated\_\_ " 0 | NMGT: Management code" "Initial Plant Growth Parameters" " 0 | IGRO: Land cover status: 0-none growing; 1-growing" ... \$ 000010003.sol: chr [1:19]" .sol file Subbasin: 2 HRU: 3 Luse: GRAS Soil: Vp38-3a-5345 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inter"| \_\_truncated\_\_ "Soil Name: Vp38-3a-5345" "Soil Hydrologic Group: C" "Maximum rooting depth(m): 910.00" ... \$ 000020000.pnd: chr [1:47]" .pnd file Subbasin: 2 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT interface" "Pond inputs:" " 0 | PND\_FR : Fraction of subbasin



area that drains into ponds. The value for PND\_FR should be between 0.0 and" | \_\_truncated\_\_ " 0  
| PND\_PSA: Surface area of ponds when filled to principal spillway [ha]" ... \$ 000020000.rte: chr  
[1:23] " .rte file Subbasin: 2 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT interface"  
"9.3638387596228 | CH\_W2 : Main channel width [m]" ".304414007785773 | CH\_D : Main chan-  
nel depth [m]" "0.03293296544532192 | CH\_S2 : Main channel slope [m/m]" ... \$ 000020000.sub:  
chr [1:64] " .sub file Subbasin: 2 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inter-  
face" "15 | SUB\_KM : Subbasin area [km2]" "" "Climate in subbasin" ... \$ 000020000.swq: chr  
[1:30] " .swq file Subbasin: 2 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inter-  
face" "Nutrient (QUAL2E parameters)" " 1 | RS1: Local algal settling rate in the reach at 20 degC  
[m/day]" " 0.05 | RS2: Benthic (sediment) source rate for dissolved phosphorus in the reach at 20  
degC [mg dissolved P/[m]" | \_\_truncated\_\_ ... \$ 000020000.wgn: chr [1:19] " .wgn file Subbasin:  
2 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT interface" " LATITUDE = 20.00  
LONGITUDE = 0.00" " ELEV [m] =2000.00" " RAIN\_YRS = 10.00" ... \$ 000020000.wus: chr  
[1:11] " .wus file Subbasin: 2 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inter-  
face" "" "" " 0.0 0.0 0.0 0.0 0.0 0.0" ... \$ 000020001.chm: chr [1:21] " .chm file Subbasin: 2 HRU:  
1 Luse: SHRB Soil: K115-2ab-5212 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow  
- SWAT inte"| \_\_truncated\_\_ "Soil Nutrient Data" " Soil Layer : 1 2 3 4 5 6 7 8 "| \_\_truncated\_\_ "  
Soil NO3 [mg/kg] : 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.00 "| \_\_truncated\_\_ ... \$ 000020001.gw  
: chr [1:14] " .gw file Subbasin: 2 HRU: 1 Luse: SHRB Soil: K115-2ab-5212 Slope: 0-5 Mon-  
day, March 21, 2011 6:57:24 PM MapWindow - SWAT inter"| \_\_truncated\_\_ " 0.5 | SHALLST  
: Initial depth of water in the shallow aquifer [mm]" " 1000 | DEEPST : Initial depth of water  
in the deep aquifer [mm]" " 31 | GW\_DELAY : Groundwater delay [days]" ... \$ 000020001.hru:  
chr [1:25] " .hru file Subbasin: 2 HRU: 1 Luse: SHRB Soil: K115-2ab-5212 Slope: 0-5 Mon-  
day, March 21, 2011 6:57:24 PM MapWindow - SWAT inte"| \_\_truncated\_\_ "0.347771714415945  
| HRU\_FR : Fraction of subbasin area contained in HRU" " 90 | SLSUBBSN : Average slope length  
[m]" "0.0260677934475378 | HRU\_SLP : Average slope stepness [m/m]" ... \$ 000020001.mgt:  
chr [1:33] " .mgt file Subbasin: 2 HRU: 1 Luse: SHRB Soil: K115-2ab-5212 Slope: 0-5 Monday,  
March 21, 2011 6:57:24 PM MapWindow - SWAT inte"| \_\_truncated\_\_ " 0 | NMGT: Management  
code" "Initial Plant Growth Parameters" " 0 | IGRO: Land cover status: 0-none growing; 1-growing"  
... \$ 000020001.sol: chr [1:19] " .sol file Subbasin: 2 HRU: 1 Luse: SHRB Soil: K115-2ab-5212  
Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inte"| \_\_truncated\_\_ "Soil  
Name: K115-2ab-5212" "Soil Hydrologic Group: C" "Maximum rooting depth(m): 580.00" ... \$  
000020002.chm: chr [1:21] " .chm file Subbasin: 2 HRU: 2 Luse: SHRB Soil: K130-2bc-4832  
Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inte"| \_\_truncated\_\_ "Soil  
Nutrient Data" " Soil Layer : 1 2 3 4 5 6 7 8 "| \_\_truncated\_\_ " Soil NO3 [mg/kg] : 0.00 0.00  
0.00 0.00 0.00 0.00 0.00 0.00 "| \_\_truncated\_\_ ... \$ 000020002.gw : chr [1:14] " .gw file Sub-  
basin: 2 HRU: 2 Luse: SHRB Soil: K130-2bc-4832 Slope: 0-5 Monday, March 21, 2011 6:57:24  
PM MapWindow - SWAT inter"| \_\_truncated\_\_ " 0.5 | SHALLST : Initial depth of water in the  
shallow aquifer [mm]" " 1000 | DEEPST : Initial depth of water in the deep aquifer [mm]" " 31  
| GW\_DELAY : Groundwater delay [days]" ... \$ 000020002.hru: chr [1:25] " .hru file Subbasin:  
2 HRU: 2 Luse: SHRB Soil: K130-2bc-4832 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM  
MapWindow - SWAT inte"| \_\_truncated\_\_ "0.351068667576475 | HRU\_FR : Fraction of subbasin  
area contained in HRU" " 90 | SLSUBBSN : Average slope length [m]" "0.0246108923185198 |  
HRU\_SLP : Average slope stepness [m/m]" ... \$ 000020002.mgt: chr [1:33] " .mgt file Subbasin:  
2 HRU: 2 Luse: SHRB Soil: K130-2bc-4832 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM  
MapWindow - SWAT inte"| \_\_truncated\_\_ " 0 | NMGT: Management code" "Initial Plant Growth  
Parameters" " 0 | IGRO: Land cover status: 0-none growing; 1-growing" ... \$ 000020002.sol: chr  
[1:19] " .sol file Subbasin: 2 HRU: 2 Luse: SHRB Soil: K130-2bc-4832 Slope: 0-5 Monday, March

21, 2011 6:57:24 PM MapWindow - SWAT inte"|\_\_truncated\_\_ "Soil Name: KI30-2bc-4832" "Soil Hydrologic Group: C" "Maximum rooting depth(m): 460.00" ... \$ 000020003.chm: chr [1:21] ".chm file Subbasin: 2 HRU: 3 Luse: GRAS Soil: Vp38-3a-5345 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inter"|\_\_truncated\_\_ "Soil Nutrient Data" " Soil Layer : 1 2 3 4 5 6 7 8 "|\_\_truncated\_\_ " Soil NO3 [mg/kg] : 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.00 "|\_\_truncated\_\_ ... \$ 000020003.gw : chr [1:14] ".gw file Subbasin: 2 HRU: 3 Luse: GRAS Soil: Vp38-3a-5345 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT interf"|\_\_truncated\_\_ " 0.5 | SHALLST : Initial depth of water in the shallow aquifer [mm]" " 1000 | DEEPST : Initial depth of water in the deep aquifer [mm]" " 31 | GW\_DELAY : Groundwater delay [days]" ... \$ 000020003.hru: chr [1:25] ".hru file Subbasin: 2 HRU: 3 Luse: GRAS Soil: Vp38-3a-5345 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inter"|\_\_truncated\_\_ "0.301159618008454 | HRU\_FR : Fraction of subbasin area contained in HRU" " 90 | SLSUBBSN : Average slope length [m]" "0.0277888825969753 | HRU\_SLP : Average slope stepness [m/m]" ... \$ 000020003.mgt: chr [1:33] ".mgt file Subbasin: 2 HRU: 3 Luse: GRAS Soil: Vp38-3a-5345 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inter"|\_\_truncated\_\_ " 0 | NMGT: Management code" "Initial Plant Growth Parameters" " 0 | IGRO: Land cover status: 0- none growing; 1-growing" ... \$ 000020003.sol: chr [1:19] ".sol file Subbasin: 2 HRU: 3 Luse: GRAS Soil: Vp38-3a-5345 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inter"|\_\_truncated\_\_ "Soil Name: Vp38-3a-5345" "Soil Hydrologic Group: C" "Maximum rooting depth(m): 910.00" ... \$ 000030000.pnd: chr [1:47] ".pnd file Subbasin: 2 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT interface" "Pond inputs:" " 0 | PND\_FR : Fraction of subbasin area that drains into ponds. The value for PND\_FR should be between 0.0 and"|\_\_truncated\_\_ " 0 | PND\_PSA: Surface area of ponds when filled to principal spillway [ha]" ... \$ 000030000.rte: chr [1:23] ".rte file Subbasin: 2 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT interface" "9.3638387596228 | CH\_W2 : Main channel width [m]" ".304414007785773 | CH\_D : Main channel depth [m]" "0.03293296544532192 | CH\_S2 : Main channel slope [m/m]" ... \$ 000030000.sub: chr [1:64] ".sub file Subbasin: 3 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT interface" "15 | SUB\_KM : Subbasin area [km2]" "" "Climate in subbasin" ... \$ 000030000.swq: chr [1:30] ".swq file Subbasin: 2 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT interface" "Nutrient (QUAL2E parameters)" " 1 | RS1: Local algal settling rate in the reach at 20 degC [m/day]" " 0.05 | RS2: Benthic (sediment) source rate for dissolved phosphorus in the reach at 20 degC [mg dissolved P/[m]"|\_\_truncated\_\_ ... \$ 000030000.wgn: chr [1:19] ".wgn file Subbasin: 2 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT interface" " LATITUDE = 20.00 LON- GITUDE = 0.00" " ELEV [m] =2000.00" " RAIN\_YRS = 10.00" ... \$ 000030000.wus: chr [1:11] ".wus file Subbasin: 2 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT interface" "" "" " 0.0 0.0 0.0 0.0 0.0 0.0" ... \$ 000030001.chm: chr [1:21] ".chm file Subbasin: 2 HRU: 1 Luse: SHRB Soil: KI15-2ab-5212 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inte"|\_\_truncated\_\_ "Soil Nutrient Data" " Soil Layer : 1 2 3 4 5 6 7 8 "|\_\_truncated\_\_ " Soil NO3 [mg/kg] : 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.00 "|\_\_truncated\_\_ ... \$ 000030001.gw : chr [1:14] ".gw file Subbasin: 2 HRU: 1 Luse: SHRB Soil: KI15-2ab-5212 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inter"|\_\_truncated\_\_ " 0.5 | SHALLST : Initial depth of water in the shallow aquifer [mm]" " 1000 | DEEPST : Initial depth of water in the deep aquifer [mm]" " 31 | GW\_DELAY : Groundwater delay [days]" ... \$ 000030001.hru: chr [1:25] ".hru file Subbasin: 2 HRU: 1 Luse: SHRB Soil: KI15-2ab-5212 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inte"|\_\_truncated\_\_ "0.347771714415945 | HRU\_FR : Fraction of subbasin area contained in HRU" " 90 | SLSUBBSN : Average slope length [m]" "0.0260677934475378 | HRU\_SLP : Average slope stepness [m/m]" ... \$ 000030001.mgt: chr [1:33] ".mgt file Subbasin: 2 HRU: 1 Luse: SHRB Soil: KI15-2ab-5212 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM

MapWindow - SWAT inte"|\_\_truncated\_\_" 0 | NMGT: Management code" "Initial Plant Growth Parameters" " 0 | IGRO: Land cover status: 0-none growing; 1-growing" ... \$ 000030001.sol: chr [1:19] ".sol file Subbasin: 2 HRU: 1 Luse: SHRB Soil: K115-2ab-5212 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inte"|\_\_truncated\_\_" "Soil Name: K115-2ab-5212" "Soil Hydrologic Group: C" "Maximum rooting depth(m): 580.00" ... \$ 000030002.chm: chr [1:21] ".chm file Subbasin: 2 HRU: 2 Luse: SHRB Soil: K130-2bc-4832 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inte"|\_\_truncated\_\_" "Soil Nutrient Data" " Soil Layer : 1 2 3 4 5 6 7 8 "|\_\_truncated\_\_" "Soil NO3 [mg/kg] : 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.00 "|\_\_truncated\_\_" ... \$ 000030002.gw : chr [1:14] ".gw file Subbasin: 2 HRU: 2 Luse: SHRB Soil: K130-2bc-4832 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inter"|\_\_truncated\_\_" 0.5 | SHALLST : Initial depth of water in the shallow aquifer [mm]" " 1000 | DEEPST : Initial depth of water in the deep aquifer [mm]" " 31 | GW\_DELAY : Groundwater delay [days]" ... \$ 000030002.hru: chr [1:25] ".hru file Subbasin: 2 HRU: 2 Luse: SHRB Soil: K130-2bc-4832 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inte"|\_\_truncated\_\_" "0.351068667576475 | HRU\_FR : Fraction of subbasin area contained in HRU" " 90 | SLSUBBSN : Average slope length [m]" "0.0246108923185198 | HRU\_SLP : Average slope stepness [m/m]" ... \$ 000030002.mgt: chr [1:33] ".mgt file Subbasin: 2 HRU: 2 Luse: SHRB Soil: K130-2bc-4832 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inte"|\_\_truncated\_\_" 0 | NMGT: Management code" "Initial Plant Growth Parameters" " 0 | IGRO: Land cover status: 0-none growing; 1-growing" ... \$ 000030002.sol: chr [1:19] ".sol file Subbasin: 2 HRU: 2 Luse: SHRB Soil: K130-2bc-4832 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inte"|\_\_truncated\_\_" "Soil Name: K130-2bc-4832" "Soil Hydrologic Group: C" "Maximum rooting depth(m): 460.00" ... \$ 000030003.chm: chr [1:21] ".chm file Subbasin: 2 HRU: 3 Luse: GRAS Soil: Vp38-3a-5345 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inter"|\_\_truncated\_\_" "Soil Nutrient Data" " Soil Layer : 1 2 3 4 5 6 7 8 "|\_\_truncated\_\_" "Soil NO3 [mg/kg] : 0.00 0.00 0.00 0.00 0.00 0.00 0.00 0.00 "|\_\_truncated\_\_" ... \$ 000030003.gw : chr [1:14] ".gw file Subbasin: 2 HRU: 3 Luse: GRAS Soil: Vp38-3a-5345 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT interf"|\_\_truncated\_\_" 0.5 | SHALLST : Initial depth of water in the shallow aquifer [mm]" " 1000 | DEEPST : Initial depth of water in the deep aquifer [mm]" " 31 | GW\_DELAY : Groundwater delay [days]" ... \$ 000030003.hru: chr [1:25] ".hru file Subbasin: 2 HRU: 3 Luse: GRAS Soil: Vp38-3a-5345 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inter"|\_\_truncated\_\_" "0.301159618008454 | HRU\_FR : Fraction of subbasin area contained in HRU" " 90 | SLSUBBSN : Average slope length [m]" "0.0277888825969753 | HRU\_SLP : Average slope stepness [m/m]" ... \$ 000030003.mgt: chr [1:33] ".mgt file Subbasin: 2 HRU: 3 Luse: GRAS Soil: Vp38-3a-5345 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inter"|\_\_truncated\_\_" 0 | NMGT: Management code" "Initial Plant Growth Parameters" " 0 | IGRO: Land cover status: 0-none growing; 1-growing" ... \$ 000030003.sol: chr [1:19] ".sol file Subbasin: 2 HRU: 3 Luse: GRAS Soil: Vp38-3a-5345 Slope: 0-5 Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT inter"|\_\_truncated\_\_" "Soil Name: Vp38-3a-5345" "Soil Hydrologic Group: C" "Maximum rooting depth(m): 910.00" ... \$ basins.bsn : chr [1:102] "Basin data .bsn file Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT interface" "Modeling Options: Land Area" "Water Balance:" " 1 | SFTMP : Snowfall temperature [ degC]" ... \$ basins.wvq : chr [1:20] "Watershed water quality file .wvq file Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT interface" " 2 | LAO : Light averaging option" " 2 | IGROPT : Algal specific growth rate option" " 50 | AIO : Ratio of chlorophyll-a to algal biomass [microg-chla/mg algae]" ... \$ crop.dat : chr [1:655] " 1 AGRL 4" " 33.50 0.45 3.00 0.15 0.05 0.50 0.95 0.64 1.00 2.00" " 30.00 11.00 0.0199 0.0032 0.0440 0.0164 0.0128 0.0060 0.0022 0.0018" " 0.250 0.2000 0.0050 4.00 0.750 8.50 660.00 36.00 0.0500 0.000" ... \$ fert.dat : chr [1:54] " 1 Elem-

```

N 1.000 0.000 0.000 0.000 0.0000.00E+000.00E+00 0.000" " 2 Elem-P 0.000 1.000 0.000 0.000
0.0000.00E+000.00E+00 0.000" " 3 ANH-NH3 0.820 0.000 0.000 0.000 1.0000.00E+000.00E+00
0.000" " 4 UREA 0.460 0.000 0.000 0.000 1.0000.00E+000.00E+00 0.000" ... $ fig.fig : chr [1:18]
"subbasin 1 1 1" " 000010000.sub" "subbasin 1 2 2" " 000020000.sub" ... $ file.cio : chr [1:71]
"Master Watershed File file.cio" "Project Description:" "General Input/Output section (file.cio):"
"Monday, March 21, 2011 6:57:24 PM MapWindow - SWAT interface" ... $ pcp1.pcp : chr
[1:14248] "Pcp" "Lati Not Used" "Long Not Used" "Elev 778" ... $ pest.dat : chr [1:233] " 1
2,4,5-TP 2600.0 0.40 5.0 20.0 0.75 2.500" " 2 2 Plus 2 20.0 0.95 10.0 21.0 0.75 660000.000" " 3
Aatrex 100.0 0.45 5.0 60.0 0.75 33.000" " 4 Abate 100000.0 0.65 5.0 30.0 0.75 0.001" ... $ till.dat
: chr [1:107] " 61 Dkplge23 0.850 100.000" " 62 Dkpllt23 0.850 100.000" " 63 Mlbr4-6 0.950
150.000" " 64 Mlbrge7 0.950 150.000" ... $ tmp1.tmp : chr [1:14248] "Tmp" "Lati Not Used"
"Long Not Used" "Elev 778" ... $ urban.dat : chr [1:18] " 1 URHD Residential-High Density 0.600
0.440" " 0.240 0.180 225.000 0.750 550.000 223.000 7.200 98.0" " 2 URMD Residential-Medium
Density 0.380 0.300" " 0.240 0.180 225.000 0.750 550.000 223.000 7.200 98.0" ...

```

## Examples

```

data(swat_general)
## maybe str(swat_general) ; plot(swat_general) ...

```

---

```

swat_objective_function

```

*A simple example objective function to be modified by user*

---

## Description

A simple example objective function to be modified by user

## Usage

```

swat_objective_function(x, calib_range, calib_params, flowgage)

```

## Arguments

x	x A vector of adjustment factors to be fed into alter_params function
calib_range	calib_range This will in the future be a range for calibration.
calib_params	See other function to make sure you have this right.
flowgage	See other function to make sure you have this right.

## Author(s)

Daniel Fuka

## Examples

```

## Not run:
Sorry, this should be custom to your project. Look and determin what you want.

## End(Not run)

```

---

 swat\_objective\_function\_rch

*An example objective function for calibrating SWAT2005 model.*


---

**Description**

An example objective function for calibrating SWAT2005 model.

**Usage**

```
swat_objective_function_rch(x, calib_range, calib_params, flowgage, rch, save_results = F)
```

**Arguments**

x	Numeric vector similar to change_params\$current defining the scalar states of the parameters to be optimized.
calib_range	2 place vector of begin and end dates for calibration.
calib_params	Dataframe in the format of change_params limited to those rows of parameters desired to calibrate. See change_params.
flowgage	list of the format of the list returned by the function get_usgs_gage.
rch	Number of the reach in the output.rch file you desire to calibrate against
save_results	Logical to express if you want to copy the optimal solution to the current directory, overwriting the swat input files.

**Value**

$\text{abs}(\text{NS} - 1)$  as needed to be able to minimize to optimal as needed for DEoptim function.

**Author(s)**

Daniel R. Fuka

---

 testSWAT2005

*A function to test the numerical correctness of the SWAT2005 exe due to bad results on some CPUs.*


---

**Description**

A function to test the numerical correctness of the SWAT2005 exe due to bad results on some CPUs.

**Usage**

```
testSWAT2005()
```

**Author(s)**

Daniel Fuka drf28@cornell.edu

---

transmissivity	<i>Transmissivity</i>
----------------	-----------------------

---

### Description

Transmissivity Fraction of direct solar radiation passing through the atmosphere based on the Bristow-Campbell eqn

### Usage

```
transmissivity(Tx,Tn, A=0.75, C=2.4, opt="1day", JD=NULL)
```

### Arguments

Tx	maximum daily temperature [C]
Tn	minimum daily temperature [C]
A	Maximum transmissivity in a location, varies with environmental conditions and elevation
C	Empirical partitioning coefficient - set to 2.4
opt	Options: "1day" uses the diurnal temperature change as Tx-Tn for any given day; "2day" depends on an average temperature change, i.e. the max temp today minus the average of the min temps of today and tomorrow; Users who do not have a continuous record of daily temperatures should use "missingdays" - note that this depends on JD input. If JD is null and "missingdays" is chosen the function will default to "1day"
JD	A vector of julian days corresponding to temp measurements. Only needed if using opt="missingdays"

### Details

Can accept either single values or vectors for Tx,Tn and JD.

### Author(s)

M. Todd Walter, Josephine Archibald

### References

Bristow KL, Campbell GS. 1984. On the relationship between incoming solar radiation and daily maximum and minimum temperature. *Agric Forest Meteorol*; 31:150-166.

**Examples**

```
## Assuming the temperature measurements are consecutive days, default option:  
transmissivity(Tx=c(10,10,10,10,10), Tn=c(1,2,6,9,2))  
## Second option:  
transmissivity(Tx=c(10,10,10,10,10), Tn=c(1,2,6,9,2), opt="2day")  
## When the days are not consecutive:  
transmissivity(Tx=c(10,10,10,10,10), Tn=c(1,2,6,9,2), JD=c(250,265,300,320,321), opt="missingdays")
```

# Index

- \*Topic **\textasciitildekwd1**
  - GroundHeat, [18](#)
- \*Topic **\textasciitildekwd2**
  - GroundHeat, [18](#)
- \*Topic **datasets**
  - change\_params, [8](#)
  - GSOD\_history, [19](#)
  - OwascoInlet, [25](#)
- \*Topic **package**
  - EcoHydRology-package, [2](#)
  
- alter\_files, [3](#)
- AtmosphericEmissivity, [4](#)
  
- BaseflowSeparation, [5](#)
- build\_gsod\_forcing\_data, [6](#)
  
- calib\_swat\_ex, [7](#)
- change\_params, [8](#)
- ConvertFlowUnits, [9](#)
  
- declination, [10](#)
  
- EcoHydRology (EcoHydRology-package), [2](#)
- EcoHydRology-package, [2](#)
- EnvirEnergy, [11](#)
- EstCloudiness, [12](#)
- EvapHeat, [13](#)
  
- get\_cfsr\_latlon, [14](#)
- get\_gsod\_stn, [15](#)
- get\_usgs\_gage, [17](#)
- GroundHeat, [18](#)
- GSOD\_history, [19](#)
  
- hydrograph, [20](#)
  
- Longwave, [21](#)
- Lumped\_VSA\_model, [22](#)
  
- NetRad, [24](#)
  
- OwascoInlet, [25](#)
  
- PET\_fromTemp, [26](#)
- PotentialSolar, [27](#)
- PotSolarInst, [28](#)
- PTpet, [29](#)
  
- RainHeat, [29](#)
- RunoffBreakdown, [30](#)
  
- SatVaporDensity, [30](#)
- SatVaporPressure, [31](#)
- SatVapPresSlope, [32](#)
- SensibleHeat, [32](#)
- setup\_swatcal, [33](#)
- slopefactor, [34](#)
- SnowMelt, [35](#)
- SoilStorage, [36](#)
- Solar, [37](#)
- solarangle, [38](#)
- SWAT2005, [38](#)
- swat\_general, [39](#)
- swat\_objective\_function, [44](#)
- swat\_objective\_function\_rch, [45](#)
  
- testSWAT2005, [45](#)
- transmissivity, [46](#)