

# Package ‘FedData’

March 12, 2018

**Type** Package

**Title** Functions to Automate Downloading Geospatial Data Available from Several Federated Data Sources

**Description** Functions to automate downloading geospatial data available from several federated data sources (mainly sources maintained by the US Federal government). Currently, the package enables extraction from seven datasets: The National Elevation Dataset digital elevation models (1 and 1/3 arc-second; USGS); The National Hydrography Dataset (USGS); The Soil Survey Geographic (SSURGO) database from the National Cooperative Soil Survey (NCSS), which is led by the Natural Resources Conservation Service (NRCS) under the USDA; the Global Historical Climatology Network (GHCN), coordinated by National Climatic Data Center at NOAA; the Daymet gridded estimates of daily weather parameters for North America, version 3, available from the Oak Ridge National Laboratory's Distributed Active Archive Center (DAAC); the International Tree Ring Data Bank; and the National Land Cover Database (NLCD).

**Version** 2.5.2

**Date** 2018-3-12

**Author** R. Kyle Bocinsky [aut, cre],  
Dylan Beaudette [ctb],  
Scott Chamberlain [ctb]

**Maintainer** R. Kyle Bocinsky <bocinsky@gmail.com>

**URL** <https://github.com/ropensci/FedData>

**BugReports** <https://github.com/ropensci/FedData/issues>

**License** MIT + file LICENSE

**Depends** R (>= 3.2.0), sp

**Imports** data.table, devtools, soilDB, igraph, curl, methods, rgdal,  
raster, Hmisc, rgeos, readr, lubridate, dplyr, magrittr,  
foreach, ncdf4, stringr, sf

**Repository** CRAN

**NeedsCompilation** no

**RoxygenNote** 6.0.1

**Suggests** testthat, covr, roxygen2

**LazyData** true

**Date/Publication** 2018-03-12 20:21:35 UTC

## R topics documented:

daymet_tiles . . . . .	2
get_daymet . . . . .	3
get_ghcn_daily . . . . .	4
get_itrdb . . . . .	8
get_ned . . . . .	10
get_nhd . . . . .	11
get_nlcd . . . . .	12
get_ssurgo . . . . .	14
nlcd_canopy_pam . . . . .	15
nlcd_impervious_pam . . . . .	15
nlcd_landcover_pam . . . . .	16
nlcd_tiles . . . . .	16
<b>Index</b>	<b>17</b>

---

daymet_tiles	<i>The DAYMET tiles SpatialPolygonsDataFrame.</i>
--------------	---

---

### Description

A dataset containing the DAYMET tiles.

### Usage

```
daymet_tiles
```

### Format

A SpatialPolygonsDataFrame with 1060 features and 5 variables:

**TileID** the numeric identifier of the tile

**XMin** the minimum longitude of the tile

**XMax** the maximum longitude of the tile

**YMin** the minimum latitude of the tile

**YMax** the maximum latitude of the tile

### Source

[https://github.com/khufkens/daymetr/blob/master/data/tile\\_outlines.rda](https://github.com/khufkens/daymetr/blob/master/data/tile_outlines.rda)

---

get_daymet	<i>Download and crop the 1-km DAYMET daily weather dataset.</i>
------------	---

---

### Description

get\_daymet returns a RasterBrick of weather data cropped to a given template study area.

### Usage

```
get_daymet(template, label, elements = NULL, years = NULL,
  raw.dir = "./RAW/DAYMET", extraction.dir = paste0("./EXTRACTIONS/", label,
  "/DAYMET"), force.redo = F)
```

### Arguments

template	A Raster* or Spatial* object to serve as a template for cropping.
label	A character string naming the study area.
elements	A character vector of elements to extract. The available elements are: dayl = Duration of the daylight period in seconds per day. This calculation is based on the period of the day during which the sun is above a hypothetical flat horizon. prcp = Daily total precipitation in millimeters per day, sum of all forms converted to water-equivalent. Precipitation occurrence on any given day may be ascertained. srad = Incident shortwave radiation flux density in watts per square meter, taken as an average over the daylight period of the day. NOTE: Daily total radiation (MJ/m2/day) can be calculated as follows: ((srad (W/m2) * dayl (s/day)) / 1,000,000) swe = Snow water equivalent in kilograms per square meter. The amount of water contained within the snowpack. tmax = Daily maximum 2-meter air temperature in degrees Celsius. tmin = Daily minimum 2-meter air temperature in degrees Celsius. vp = Water vapor pressure in pascals. Daily average partial pressure of water vapor.
years	A numeric vector of years to extract.
raw.dir	A character string indicating where raw downloaded files should be put. The directory will be created if missing. Defaults to './RAW/DAYMET/'.
extraction.dir	A character string indicating where the extracted and cropped DEM should be put. The directory will be created if missing. Defaults to './EXTRACTIONS/DAYMET/'.
force.redo	If an extraction for this template and label already exists, should a new one be created?

**Value**

A named list of RasterBricks of weather data cropped to the extent of the template.

**Examples**

```
## Not run:
# Extract data for the Village Ecodynamics Project 'VEPIIN' study area:
# http://village.anth.wsu.edu
vepPolygon <- polygon_from_extent(raster::extent(672800,740000,4102000,4170000),
  proj4string='+proj=utm +datum=NAD83 +zone=12')

# Get the DAYMET (North America only)
# Returns a list of raster bricks
DAYMET <- get_daymet(template=vepPolygon,
  label='VEPIIN',
  elements = c('prcp','tmin','tmax'),
  years = 1980:1985)

# Plot with raster::plot
plot(DAYMET$tmin$X1985.10.23)

## End(Not run)
```

---

get_ghcn_daily	<i>Download and crop the Global Historical Climate Network-Daily data.</i>
----------------	--

---

**Description**

get\_ghcn\_daily returns a named list of length 2:

1. 'spatial': A SpatialPointsDataFrame of the locations of GHCN weather stations in the template, and
2. 'tabular': A named list of [data.frames](#) with the daily weather data for each station. The name of each list item is the station ID.

**Usage**

```
get_ghcn_daily(template = NULL, label = NULL, elements = NULL,
  years = NULL, raw.dir = "./RAW/GHCN",
  extraction.dir = paste0("./EXTRACTIONS/", label, "/GHCN"),
  standardize = F, force.redo = F)
```

**Arguments**

template	A Raster* or Spatial* object to serve as a template for cropping. Alternatively, a character vector providing GHCN station IDs. If missing, all stations will be downloaded!
----------	--

label	A character string naming the study area.
elements	<p>A character vector of elements to extract.</p> <p>The five core elements are:</p> <p>PRCP = Precipitation (tenths of mm)</p> <p>SNOW = Snowfall (mm)</p> <p>SNWD = Snow depth (mm)</p> <p>TMAX = Maximum temperature (tenths of degrees C)</p> <p>TMIN = Minimum temperature (tenths of degrees C)</p> <p>The other elements are:</p> <p>ACMC = Average cloudiness midnight to midnight from 30-second ceilometer data (percent)</p> <p>ACMH = Average cloudiness midnight to midnight from manual observations (percent)</p> <p>ACSC = Average cloudiness sunrise to sunset from 30-second ceilometer data (percent)</p> <p>ACSH = Average cloudiness sunrise to sunset from manual observations (percent)</p> <p>AWDR = Average daily wind direction (degrees)</p> <p>AWND = Average daily wind speed (tenths of meters per second)</p> <p>DAEV = Number of days included in the multiday evaporation total (MDEV)</p> <p>DAPR = Number of days included in the multiday precipitation total (MDPR)</p> <p>DASF = Number of days included in the multiday snowfall total (MDSF)</p> <p>DATN = Number of days included in the multiday minimum temperature (MDTN)</p> <p>DATX = Number of days included in the multiday maximum temperature (MDTX)</p> <p>DAWM = Number of days included in the multiday wind movement (MDWM)</p> <p>DWPR = Number of days with non-zero precipitation included in multiday precipitation total (MDPR)</p> <p>EVAP = Evaporation of water from evaporation pan (tenths of mm)</p> <p>FMTM = Time of fastest mile or fastest 1-minute wind (hours and minutes, i.e., HHMM)</p> <p>FRGB = Base of frozen ground layer (cm)</p> <p>FRGT = Top of frozen ground layer (cm)</p> <p>FRTH = Thickness of frozen ground layer (cm)</p> <p>GAHT = Difference between river and gauge height (cm)</p> <p>MDEV = Multiday evaporation total (tenths of mm; use with DAEV)</p> <p>MDPR = Multiday precipitation total (tenths of mm; use with DAPR and DWPR, if available)</p> <p>MDSF = Multiday snowfall total</p> <p>MDTN = Multiday minimum temperature (tenths of degrees C; use with DATN)</p> <p>MDTX = Multiday maximum temperature (tenths of degrees C; use with DATX)</p> <p>MDWM = Multiday wind movement (km)</p> <p>MNPN = Daily minimum temperature of water in an evaporation pan (tenths of degrees C)</p> <p>MXPN = Daily maximum temperature of water in an evaporation pan (tenths of degrees C)</p> <p>PGTM = Peak gust time (hours and minutes, i.e., HHMM)</p>

PSUN = Daily percent of possible sunshine (percent)

SN\*# = Minimum soil temperature (tenths of degrees C) where \* corresponds to a code for ground cover and # corresponds to a code for soil depth.

Ground cover codes include the following:

0 = unknown

1 = grass

2 = fallow

3 = bare ground

4 = brome grass

5 = sod

6 = straw mulch

7 = grass muck

8 = bare muck

Depth codes include the following:

1 = 5 cm

2 = 10 cm

3 = 20 cm

4 = 50 cm

5 = 100 cm

6 = 150 cm

7 = 180 cm

SX\*# = Maximum soil temperature (tenths of degrees C) where \* corresponds to a code for ground cover and # corresponds to a code for soil depth.

See SN\*# for ground cover and depth codes.

TAVG = Average temperature (tenths of degrees C) [Note that TAVG from source 'S' corresponds to an average for the period ending at 2400 UTC rather than local midnight]

THIC = Thickness of ice on water (tenths of mm)

TOBS = Temperature at the time of observation (tenths of degrees C)

TSUN = Daily total sunshine (minutes)

WDF1 = Direction of fastest 1-minute wind (degrees)

WDF2 = Direction of fastest 2-minute wind (degrees)

WDF5 = Direction of fastest 5-second wind (degrees)

WDFG = Direction of peak wind gust (degrees)

WDFI = Direction of highest instantaneous wind (degrees)

WDFM = Fastest mile wind direction (degrees)

WDMV = 24-hour wind movement (km)

WESD = Water equivalent of snow on the ground (tenths of mm)

WESF = Water equivalent of snowfall (tenths of mm)

WSF1 = Fastest 1-minute wind speed (tenths of meters per second)

WSF2 = Fastest 2-minute wind speed (tenths of meters per second)

WSF5 = Fastest 5-second wind speed (tenths of meters per second)

WSFG = Peak gust wind speed (tenths of meters per second)

WSFI = Highest instantaneous wind speed (tenths of meters per second)

WSFM = Fastest mile wind speed (tenths of meters per second)

WT\*\* = Weather Type where \*\* has one of the following values:

01 = Fog, ice fog, or freezing fog (may include heavy fog)  
 02 = Heavy fog or heaving freezing fog (not always distinguished from fog)  
 03 = Thunder  
 04 = Ice pellets, sleet, snow pellets, or small hail  
 05 = Hail (may include small hail)  
 06 = Glaze or rime  
 07 = Dust, volcanic ash, blowing dust, blowing sand, or blowing obstruction  
 08 = Smoke or haze  
 09 = Blowing or drifting snow  
 10 = Tornado, waterspout, or funnel cloud  
 11 = High or damaging winds  
 12 = Blowing spray  
 13 = Mist  
 14 = Drizzle  
 15 = Freezing drizzle  
 16 = Rain (may include freezing rain, drizzle, and freezing drizzle)  
 17 = Freezing rain  
 18 = Snow, snow pellets, snow grains, or ice crystals  
 19 = Unknown source of precipitation  
 21 = Ground fog  
 22 = Ice fog or freezing fog

WV\*\* = Weather in the Vicinity where \*\* has one of the following values:

01 = Fog, ice fog, or freezing fog (may include heavy fog)  
 03 = Thunder  
 07 = Ash, dust, sand, or other blowing obstruction  
 18 = Snow or ice crystals  
 20 = Rain or snow shower

years            A numeric vector indicating which years to get.  
 raw.dir         A character string indicating where raw downloaded files should be put. The directory will be created if missing. Defaults to './RAW/GHCN/'.  
 extraction.dir   A character string indicating where the extracted and cropped GHCN shapefiles should be put. The directory will be created if missing. Defaults to './EXTRACTIONS/GHCN/'.  
 standardize     Select only common year/month/day? Defaults to FALSE.  
 force.redo       If an extraction for this template and label already exists, should a new one be created? Defaults to FALSE.

### Value

A named list containing the 'spatial' and 'tabular' data.

### Examples

```
## Not run:
# Extract data for the Village Ecodynamics Project 'VEPIIN' study area:
```

```

# http://village.anth.wsu.edu
vepPolygon <- polygon_from_extent(raster::extent(672800,740000,4102000,4170000),
  proj4string='+proj=utm +datum=NAD83 +zone=12')

# Get the daily GHCN data (GLOBAL)
# Returns a list: the first element is the spatial locations of stations,
# and the second is a list of the stations and their daily data
GHCN.prcp <- get_ghcn_daily(template=vepPolygon, label='VEPIIN', elements=c('prcp'))

# Plot the VEP polygon
plot(vepPolygon)

# Plot the spatial locations
plot(GHCN.prcp$spatial, pch=1, add=T)
legend('bottomleft', pch=1, legend='GHCN Precipitation Records')

# Elements for which you require the same data
# (i.e., minimum and maximum temperature for the same days)
# can be standardized using standardize==T
GHCN.temp <- get_ghcn_daily(template=vepPolygon,
  label='VEPIIN',
  elements=c('tmin','tmax'),
  standardize=T)

# Plot the VEP polygon
plot(vepPolygon)

# Plot the spatial locations
plot(GHCN.temp$spatial, pch=1, add=T)
legend('bottomleft', pch=1, legend='GHCN Temperature Records')

## End(Not run)

```

---

get\_itrdb

*Download the latest version of the ITRDB, and extract given parameters.*

---

## Description

get\_itrdb returns a named list of length 3:

1. 'metadata': A data.table or SpatialPointsDataFrame (if makeSpatial==TRUE) of the locations and names of extracted ITRDB chronologies,
2. 'widths': A matrix of tree-ring widths/densities given user selection, and
3. 'depths': A matrix of tree-ring sample depths.



**Usage**

```
get_itrdb(template = NULL, label = NULL, recon.years = NULL,
  calib.years = NULL, species = NULL, measurement.type = NULL,
  chronology.type = NULL, makeSpatial = F, raw.dir = "./RAW/ITRDB",
  extraction.dir = ifelse(!is.null(label), paste0("./EXTRACTIONS/", label,
  "/ITRDB"), "./EXTRACTIONS/ITRDB"), force.redo = FALSE)
```

**Arguments**

template	A Raster* or Spatial* object to serve as a template for selecting chronologies. If missing, all available global chronologies are returned.
label	A character string naming the study area.
recon.years	A numeric vector of years over which reconstructions are needed; if missing, the union of all years in the available chronologies are given.
calib.years	A numeric vector of all required years—chronologies without these years will be discarded; if missing, all available chronologies are given.
species	A character vector of 4-letter tree species identifiers; if missing, all available chronologies are given.
measurement.type	A character vector of measurement type identifiers. Options include: <ul style="list-style-type: none"> <li>• 'Total Ring Density'</li> <li>• 'Earlywood Width'</li> <li>• 'Earlywood Density'</li> <li>• 'Latewood Width'</li> <li>• 'Minimum Density'</li> <li>• 'Ring Width'</li> <li>• 'Latewood Density'</li> <li>• 'Maximum Density'</li> <li>• 'Latewood Percent'</li> </ul> if missing, all available chronologies are given.
chronology.type	A character vector of chronology type identifiers. Options include: <ul style="list-style-type: none"> <li>• 'ARSTND'</li> <li>• 'Low Pass Filter'</li> <li>• 'Residual'</li> <li>• 'Standard'</li> <li>• 'Re-Whitened Residual'</li> <li>• 'Measurements Only'</li> </ul> if missing, all available chronologies are given.
makeSpatial	Should the metadata be presented as a SpatialPointsDataFrame? Defaults to FALSE.
raw.dir	A character string indicating where raw downloaded files should be put. The directory will be created if missing. Defaults to './RAW/ITRDB/'.

`extraction.dir` A character string indicating where the extracted and cropped ITRDB dataset should be put. The directory will be created if missing. Defaults to `./EXTRACTIONS/ITRDB/`.

`force.redo` If an extraction already exists, should a new one be created? Defaults to `FALSE`.

### Value

A named list containing the `'metadata'`, `'widths'`, and `'depths'` data.

### Examples

```
## Not run:
# Extract data for the Village Ecodynamics Project 'VEPIIN' study area:
# http://village.anth.wsu.edu
vepPolygon <- polygon_from_extent(raster::extent(672800,740000,4102000,4170000),
  proj4string='+proj=utm +datum=NAD83 +zone=12')

# Get the ITRDB records
ITRDB <- get_itrdb(template=vepPolygon, label='VEPIIN', makeSpatial=T)

# Plot the VEP polygon
plot(vepPolygon)

# Map the locations of the tree ring chronologies
plot(ITRDB$metadata, pch=1, add=T)
legend('bottomleft', pch=1, legend='ITRDB chronologies')

## End(Not run)
```

---

<code>get_ned</code>	<i>Download and crop the 1 (~30 meter) or 1/3 (~10 meter) arc-second National Elevation Dataset.</i>
----------------------	--

---

### Description

`get_ned` returns a `RasterLayer` of elevation data cropped to a given template study area.

### Usage

```
get_ned(template, label, res = "1", raw.dir = "./RAW/NED",
  extraction.dir = paste0("./EXTRACTIONS/", label, "/NED"),
  raster.options = c("COMPRESS=DEFLATE", "ZLEVEL=9", "INTERLEAVE=BAND"),
  force.redo = F)
```

**Arguments**

template	A Raster* or Spatial* object to serve as a template for cropping.
label	A character string naming the study area.
res	A character string representing the desired resolution of the NED. '1' indicates the 1 arc-second NED (the default), while '13' indicates the 1/3 arc-second dataset.
raw.dir	A character string indicating where raw downloaded files should be put. The directory will be created if missing. Defaults to './RAW/NED/'.
extraction.dir	A character string indicating where the extracted and cropped DEM should be put. The directory will be created if missing. Defaults to './EXTRACTIONS/NED/'.
raster.options	a vector of options for raster::writeRaster.
force.redo	If an extraction for this template and label already exists, should a new one be created?

**Value**

A RasterLayer DEM cropped to the extent of the template.

**Examples**

```
## Not run:
# Extract data for the Village Ecodynamics Project 'VEPIIN' study area:
# http://village.anth.wsu.edu
vepPolygon <- polygon_from_extent(raster::extent(672800,740000,4102000,4170000),
  proj4string='+proj=utm +datum=NAD83 +zone=12')

# Get the NED (USA ONLY)
# Returns a raster
NED <- get_ned(template=vepPolygon, label='VEPIIN')

# Plot with raster::plot
plot(NED)

## End(Not run)
```

---

get\_nhd

*Download and crop the National Hydrography Dataset.*


---

**Description**

get\_nhd returns a list of Spatial\* objects extracted from the National Hydrography Dataset.

**Usage**

```
get_nhd(template, label, raw.dir = "./RAW/NHD",
  extraction.dir = paste0("./EXTRACTIONS/", label, "/NHD"),
  force.redo = FALSE)
```

**Arguments**

template	A Raster* or Spatial* object to serve as a template for cropping.
label	A character string naming the study area.
raw.dir	A character string indicating where raw downloaded files should be put. The directory will be created if missing. Defaults to './RAW/NHD/'.
extraction.dir	A character string indicating where the extracted and cropped NHD shapefiles should be put. The directory will be created if missing. Defaults to './EXTRACTIONS/NHD/'.
force.redo	If an extraction for this template and label already exists, should a new one be created?

**Value**

A list of Spatial\* objects extracted from the National Hydrography Dataset.

**Examples**

```
## Not run:
# Extract data for the Village Ecodynamics Project 'VEPIIN' study area:
# http://village.anth.wsu.edu
vepPolygon <- polygon_from_extent(raster::extent(672800,740000,4102000,4170000),
  proj4string='+proj=utm +datum=NAD83 +zone=12')

# Get the NHD (USA ONLY)
NHD <- get_nhd(template=vepPolygon, label='VEPIIN')

# Plot the VEP polygon
plot(vepPolygon)

# Plot the NHD data
plot(NHD$NHDFlowline, add=T)
plot(NHD$NHDLine, add=T)
plot(NHD$NHDArea, col='black', add=T)
plot(NHD$NHDWaterbody, col='black', add=T)

## End(Not run)
```

---

get\_nlcd

*Download and crop the National Land Cover Database.*


---

**Description**

get\_nlcd returns a RasterLayer of NLCD data cropped to a given template study area.

**Usage**

```
get_nlcd(template, label, year = 2011, dataset = "landcover",
  raw.dir = "./RAW/NLCD", extraction.dir = paste0("./EXTRACTIONS/", label,
  "/NLCD"), raster.options = c("COMPRESS=DEFLATE", "ZLEVEL=9",
  "INTERLEAVE=BAND"), force.redo = F)
```

**Arguments**

template	An Raster* or Spatial* object to serve as a template for cropping.
label	A character string naming the study area.
year	An integer representing the year of desired NLCD product. Acceptable values are 2011 (default), 2006, and 2001.
dataset	A character string representing type of the NLCD product. Acceptable values are 'landcover' (default), 'impervious', and 'canopy'. As of February 7, 2018, the canopy data for 2006 are not available through the National Map Staged datasets, and so aren't available in FedData.
raw.dir	A character string indicating where raw downloaded files should be put. The directory will be created if missing. Defaults to './RAW/NLCD/'.
extraction.dir	A character string indicating where the extracted and cropped DEM should be put. The directory will be created if missing. Defaults to './EXTRACTIONS/NLCD/'.
raster.options	a vector of options for raster::writeRaster.
force.redo	If an extraction for this template and label already exists, should a new one be created?

**Value**

A RasterLayer DEM cropped to the extent of the template.

**Examples**

```
## Not run:
# Extract data for the Village Ecodynamics Project 'VEPIIN' study area:
# http://village.anth.wsu.edu
vepPolygon <- polygon_from_extent(raster::extent(672800, 740000, 4102000, 4170000),
  proj4string='+proj=utm +datum=NAD83 +zone=12')

# Get the NLCD (USA ONLY)
# Returns a raster
NLCD <- get_nlcd(template=vepPolygon, label='VEPIIN')

# Plot with raster::plot
plot(NLCD)

## End(Not run)
```

---

 get\_ssurgo

*Download and crop data from the NRCS SSURGO soils database.*


---

### Description

This is an efficient method for spatially merging several different soil survey areas as well as merging their tabular data.

### Usage

```
get_ssurgo(template, label, raw.dir = "./RAW/SSURGO",
           extraction.dir = paste0("./EXTRACTIONS/", label, "/SSURGO"),
           force.redo = FALSE)
```

### Arguments

template	A Raster* or Spatial* object to serve as a template for cropping; optionally, a vector of area names [e.g., c('IN087','IN088')] may be provided.
label	A character string naming the study area.
raw.dir	A character string indicating where raw downloaded files should be put. The directory will be created if missing. Defaults to './RAW/SSURGO/'.
extraction.dir	A character string indicating where the extracted and cropped SSURGO shapefiles should be put. The directory will be created if missing. Defaults to './EXTRACTIONS/SSURGO/'.
force.redo	If an extraction for this template and label already exists, should a new one be created? Defaults to FALSE.

### Details

get\_ssurgo returns a named list of length 2:

1. 'spatial': A SpatialPolygonsDataFrame of soil mapunits in the template, and
2. 'tabular': A named list of [data.frames](#) with the SSURGO tabular data.

### Value

A named list containing the 'spatial' and 'tabular' data.

### Examples

```
## Not run:
# Extract data for the Village Ecodynamics Project 'VEPIIN' study area:
# http://village.anth.wsu.edu
vepPolygon <- polygon_from_extent(raster::extent(672800,740000,4102000,4170000),
                                proj4string='+proj=utm +datum=NAD83 +zone=12')

# Get the NRCS SSURGO data (USA ONLY)
```

```
SSURGO.VEPIIN <- get_ssurgo(template=vepPolygon, label='VEPIIN')

# Plot the VEP polygon
plot(vepPolygon)

# Plot the SSURGO mapunit polygons
plot(SSURGO.VEPIIN$spatial, lwd=0.1, add=T)

# Or, download by Soil Survey Area names
SSURGO.areas <- get_ssurgo(template=c('C0670', 'C0075'), label='CO_TEST')

# Let's just look at spatial data for C0675
SSURGO.areas.C0675 <- SSURGO.areas$spatial[SSURGO.areas$spatial$AREASYMBOL=='C0075',]

# And get the NED data under them for pretty plotting
NED.C0675 <- get_ned(template=SSURGO.areas.C0675, label='SSURGO_C0675')

# Plot the SSURGO mapunit polygons, but only for C0675
plot(NED.C0675)
plot(SSURGO.areas.C0675, lwd=0.1, add=T)

## End(Not run)
```

---

nlcd\_canopy\_pam

*The NLCD canopy PAM attributes.*

---

### **Description**

A dataset containing the PAM attributes.

### **Usage**

```
nlcd_canopy_pam
```

### **Format**

An object of class character of length 2345.

---

nlcd\_impervious\_pam

*The NLCD impervious PAM attributes.*

---

### **Description**

A dataset containing the PAM attributes.

### **Usage**

```
nlcd_impervious_pam
```

**Format**

An object of class character of length 2345.

---

nlcd\_landcover\_pam      *The NLCD landcover PAM attributes.*

---

**Description**

A dataset containing the PAM attributes.

**Usage**

```
nlcd_landcover_pam
```

**Format**

An object of class character of length 2606.

---

nlcd\_tiles              *The NLCD tiles SpatialPolygonsDataFrame.*

---

**Description**

A dataset containing the NLCD tiles.

**Usage**

```
nlcd_tiles
```

**Format**

A SpatialPolygonsDataFrame with 203 features and 1 variable:

**Name** the name of the tile



# Index

## \*Topic **datasets**

- daymet\_tiles, [2](#)
- nlcd\_canopy\_pam, [15](#)
- nlcd\_impervious\_pam, [15](#)
- nlcd\_landcover\_pam, [16](#)
- nlcd\_tiles, [16](#)

- data.frame, [4](#), [14](#)
- daymet\_tiles, [2](#)

- get\_daymet, [3](#)
- get\_ghcn\_daily, [4](#)
- get\_itrdb, [8](#)
- get\_ned, [10](#)
- get\_nhd, [11](#)
- get\_nlcd, [12](#)
- get\_ssurgo, [14](#)

- nlcd\_canopy\_pam, [15](#)
- nlcd\_impervious\_pam, [15](#)
- nlcd\_landcover\_pam, [16](#)
- nlcd\_tiles, [16](#)