

Package ‘MittagLeffleR’

September 6, 2021

Type Package

Title Mittag-Leffler Family of Distributions

Version 0.4.1

Author Katharina Hees [aut],
Peter Straka [aut, cre],
Gurtek Gill [aut],
Roberto Garrappa [ctb]

Maintainer Peter Straka <straka.ps@gmail.com>

Description Implements the Mittag-Leffler function, distribution,
random variate generation, and estimation. Based on the Laplace-Inversion
algorithm by Garrappa, R. (2015) <doi:10.1137/140971191>.

URL <https://strakaps.github.io/MittagLeffleR/>

BugReports <https://github.com/strakaps/MittagLeffleR/issues>

License GPL (>= 2)

Encoding UTF-8

RoxygenNote 7.1.1

Imports stats, stabledist

Suggests knitr, rmarkdown, testthat, animation, magrittr

VignetteBuilder knitr

NeedsCompilation no

Repository CRAN

Date/Publication 2021-09-06 06:50:02 UTC

R topics documented:

MittagLeffleR-package	2
logMomentEstimator	3
Mittag-Leffler	3
mlf	5
mlmle	6
Index	7

MittagLeffleR-package *Mittag-Leffler family of distributions*

Description

A generalization of the exponential distribution. Contains

- the Mittag-Leffler function `mlf`
- distributions (`dml`, `pml`, `qml`) and random variate generation (`rml`)
- a log-moment estimator (`logMomentEstimator`), and maximum likelihood estimator (`mlmle`)

Details

- [Plots of the Mittag-Leffler distributions](#)
- [Details of Mittag-Leffler random variate generation](#)
- [Probabilities and Quantiles](#)

Also see the package web page at <https://strakaps.github.io/MittagLeffleR/reference/index.html>

Author(s)

Maintainer: Peter Straka <straka.ps@gmail.com>

Authors:

- Katharina Hees <heeskatharina@gmail.com>
- Gurtek Gill

Other contributors:

- Roberto Garrappa <roberto.garrappa@uniba.it> [contributor]

See Also

Useful links:

- <https://strakaps.github.io/MittagLeffleR/>
- Report bugs at <https://github.com/strakaps/MittagLeffleR/issues>

logMomentEstimator *Log-Moments Estimator for the Mittag-Leffler Distribution (Type 1).*

Description

Tail and scale parameter of the Mittag-Leffler distribution are estimated by matching with the first two empirical log-moments (see Cahoy et al., doi: [10.1016/j.jspi.2010.04.016](https://doi.org/10.1016/j.jspi.2010.04.016)).

Usage

```
logMomentEstimator(x, alpha = 0.05)
```

Arguments

x	A vector of non-negative data.
alpha	Confidence intervals are calculated at level 1 - alpha.

Value

A named vector with entries (nu, delta, nuLo, nuHi, deltaLo, deltaHi) where nu is the tail parameter and delta the scale parameter of the Mittag-Leffler distribution, with confidence intervals (nuLo, nuHi) resp. (deltaLo, deltaHi).

References

Cahoy, D. O., Uchaikin, V. V., & Woyczynski Wojbor, W. A. (2010). Parameter estimation for fractional Poisson processes. *Journal of Statistical Planning and Inference*, 140(11), 3106–3120. doi: [10.1016/j.jspi.2010.04.016](https://doi.org/10.1016/j.jspi.2010.04.016)

Cahoy, D. O. (2013). Estimation of Mittag-Leffler Parameters. *Communications in Statistics - Simulation and Computation*, 42(2), 303–315. doi: [10.1080/03610918.2011.640094](https://doi.org/10.1080/03610918.2011.640094)

Examples

```
logMomentEstimator(rml(n = 1000, scale = 0.03, tail = 0.84), alpha=0.95)
```

Mittag-Leffler *Distribution functions and random number generation.*

Description

Probability density, cumulative distribution function, quantile function and random variate generation for the two types of Mittag-Leffler distribution. The Laplace inversion algorithm by Garrappa is used for the pdf and cdf (see <https://www.mathworks.com/matlabcentral/fileexchange/48154-the-mittag-leffler-function>).

Usage

```
dml(x, tail, scale = 1, log = FALSE, second.type = FALSE)
```

```
pm1(q, tail, scale = 1, second.type = FALSE, lower.tail = TRUE, log.p = FALSE)
```

```
qm1(p, tail, scale = 1, second.type = FALSE, lower.tail = TRUE, log.p = FALSE)
```

```
rml(n, tail, scale = 1, second.type = FALSE)
```

Arguments

<code>x, q</code>	vector of quantiles.
<code>tail</code>	tail parameter.
<code>scale</code>	scale parameter.
<code>log, log.p</code>	logical; if TRUE, probabilities <code>p</code> are given as $\log(p)$.
<code>second.type</code>	logical; if FALSE (default), first type of Mittag-Leffler distribution is assumed.
<code>lower.tail</code>	logical; if TRUE, probabilities are $P[X \leq x]$ otherwise, $P[X > x]$
<code>p</code>	vector of probabilities.
<code>n</code>	number of random draws.

Details

The Mittag-Leffler function `mlf` defines two types of probability distributions:

The **first type** of Mittag-Leffler distribution assumes the Mittag-Leffler function as its tail function, so that the CDF is given by

$$F(q; \alpha, \tau) = 1 - E_{\alpha,1}(-(q/\tau)^\alpha)$$

for $q \geq 0$, tail parameter $0 < \alpha \leq 1$, and scale parameter $\tau > 0$. Its PDF is given by

$$f(x; \alpha, \tau) = x^{\alpha-1} E_{\alpha,\alpha}[-(x/\tau)^\alpha] / \tau^\alpha.$$

As α approaches 1 from below, the Mittag-Leffler converges (weakly) to the exponential distribution. For $0 < \alpha < 1$, it is (very) heavy-tailed, i.e. has infinite mean.

The **second type** of Mittag-Leffler distribution is defined via the Laplace transform of its density `f`:

$$\int_0^\infty \exp(-sx) f(x; \alpha, 1) dx = E_{\alpha,1}(-s)$$

It is light-tailed, i.e. all its moments are finite. At scale τ , its density is

$$f(x; \alpha, \tau) = f(x/\tau; \alpha, 1) / \tau.$$

Value

`dml` returns the density, `pm1` returns the distribution function, `qm1` returns the quantile function, and `rml` generates random variables.

References

Haubold, H. J., Mathai, A. M., & Saxena, R. K. (2011). Mittag-Leffler Functions and Their Applications. *Journal of Applied Mathematics*, 2011, 1–51. doi: [10.1155/2011/298628](https://doi.org/10.1155/2011/298628)

Mittag-Leffler distribution. (2017, May 3). In Wikipedia, The Free Encyclopedia. https://en.wikipedia.org/w/index.php?title=Mittag-Leffler_distribution&oldid=778429885

Examples

```
dml(1, 0.8)
dml(1, 0.6, second.type=TRUE)
pml(2, 0.7, 1.5)
qml(p = c(0.25, 0.5, 0.75), tail = 0.6, scale = 100)
rml(10, 0.7, 1)
```

 mlf

Mittag-Leffler Function.

Description

The generalized (two-parameter) Mittag-Leffler function is defined by the power series

$$E_{\alpha,\beta}(z) = \sum_{k=0}^{\infty} z^k / \Gamma(\alpha k + \beta)$$

for complex z and complex α, β with $Real(\alpha) > 0$ (only implemented for real valued parameters).

Usage

```
mlf(z, a, b = 1, g = 1)
```

Arguments

<code>z</code>	The argument (real-valued)
<code>a, b, g</code>	Parameters of the Mittag-Leffler distribution; see Garrappa

Value

mlf returns the value of the Mittag-Leffler function.

References

Garrappa, R. (2015). Numerical Evaluation of Two and Three Parameter Mittag-Leffler Functions. *SIAM Journal on Numerical Analysis*, 53(3), 1350–1369. doi: [10.1137/140971191](https://doi.org/10.1137/140971191)

The Mittag-Leffler function. MathWorks File Exchange. <https://au.mathworks.com/matlabcentral/fileexchange/48154-the-mittag-leffler-function>

Examples

```
mlf(2,0.7)
```

mlmle

Maximum Likelihood Estimation of the Mittag-Leffler distribution

Description

Optimizes the bivariate loglikelihood of the Mittag-Leffler distribution via [optim](#). Uses [logMomentEstimator](#) for initial parameter values.

Usage

```
mlmle(data, ...)
```

Arguments

data	Vector of class "numeric"
...	Additional parameters passed on to optim .

Value

The output of [optim](#).

Examples

```
library(magrittr)
rml(n = 100, tail = 0.8, scale = 1000) %>% mlmle()
```

Index

* Mittag Leffler Distribution

Mittag-Leffler, 3

dml, 2

dml (Mittag-Leffler), 3

logMomentEstimator, 2, 3, 6

Mittag-Leffler, 3

MittagLeffleR (MittagLeffleR-package), 2

MittagLeffleR-package, 2

mlf, 2, 4, 5

mlmle, 2, 6

optim, 6

pml, 2

pml (Mittag-Leffler), 3

qml, 2

qml (Mittag-Leffler), 3

rml, 2

rml (Mittag-Leffler), 3