

Package ‘Qapprox’

September 29, 2020

Type Package

Title Approximation to the Survival Functions of Quadratic Forms of Gaussian Variables

Version 0.1.0

Author Hong Zhang

Maintainer Hong Zhang <hzhang@wpi.edu>

Description Calculates the right-tail probability of quadratic forms of Gaussian variables using the skewness-kurtosis ratio matching method, modified Liu-Tang-Zhang method and Satterthwaite-Welch method. The technical details can be found in Hong Zhang, Judong Shen and Zheyang Wu (2020) <arXiv:2005.00905>.

License GPL-2

Imports stats

Encoding UTF-8

LazyData true

RoxygenNote 6.1.0

NeedsCompilation no

Repository CRAN

Date/Publication 2020-09-29 08:10:06 UTC

R topics documented:

Qapprox	2
Qapprox_nc	3

Index	4
-------	---

Qapprox	<i>Right-tail probability of quadratic forms of centered Gaussian variables.</i>
---------	--

Description

Right-tail probability of quadratic forms of centered Gaussian variables.

Usage

```
Qapprox(q, Sigma, A = NULL, method = "MR")
```

Arguments

q	- quantile, could be a vector.
Sigma	- covariance matrix of Gaussian variables.
A	- a positive-semi-definite matrix that defines the quadratic form.
method	- "MR": moment-ratio (skewness-kurtosis) matching method; "SW": Satterthwaite-Welch method that matches mean and variance; "LTZ4": Liu-Tang-Zhang method that matches the kurtosis.

Value

The right-tail probability of a quadratic form ($Q = X'AX$) of centered Gaussian variables.

References

1. Hong Zhang, Judong Shen and Zheyang Wu. "An efficient and accurate approximation to the distribution of quadratic forms of Gaussian variables", arXiv:2005.00905.

Examples

```
n = 100
Sigma = toeplitz(1/(1:n))
thr = 180
Qapprox(thr, Sigma, method="SW")
Qapprox(thr, Sigma, method="LTZ4")
Qapprox(thr, Sigma, method="MR")
```

Qapprox_nc	<i>Right-tail probability of quadratic forms ($Q = X'AX$) of noncentral Gaussian variables.</i>
------------	--

Description

Right-tail probability of quadratic forms ($Q = X'AX$) of noncentral Gaussian variables.

Usage

```
Qapprox_nc(q, mu, Sigma, A = NULL, method = "MR")
```

Arguments

q	- quantile, could be a vector.
mu	- mean vector of Gaussian variables.
Sigma	- covariance matrix of Gaussian variables.
A	- a positive-semi-definite matrix that defines the quadratic form.
method	- "MR": moment-ratio (skewness-kurtosis) matching method; "SW": Satterthwaite-Welch method that matches mean and variance; "LTZ4": Liu-Tang-Zhang method that matches the kurtosis.

Value

The right-tail probability of a quadratic form ($Q = X'AX$) of noncentral Gaussian variables.

References

1. Hong Zhang, Judong Shen and Zheyang Wu. "An efficient and accurate approximation to the distribution of quadratic forms of Gaussian variables", arXiv:2005.00905.

Examples

```
n = 100
Sigma = toeplitz(1/(1:n))
mu = rep(1, n)
thr = 500
Qapprox_nc(thr, mu, Sigma, method="SW")
Qapprox_nc(thr, mu, Sigma, method="LTZ4")
Qapprox_nc(thr, mu, Sigma, method="MR")
```

Index

Qapprox, [2](#)
Qapprox_nc, [3](#)