

Package ‘StreamMetabolism’

October 12, 2022

Type Package

Title Calculate Single Station Metabolism from Diurnal Oxygen Curves

Version 1.1.2

Date 2016-08-24

Author Stephen A Sefick Jr.

Maintainer Stephen A Sefick Jr. <ssefick@gmail.com>

URL <https://github.com/ssefick/StreamMetabolism>

Depends zoo, chron, maptools

Description I provide functions to calculate Gross Primary Productivity, Net Ecosystem Production, and Ecosystem Respiration from single station diurnal Oxygen curves.

License GPL (>= 3)

LazyLoad yes

Repository CRAN

NeedsCompilation no

Date/Publication 2016-09-12 20:55:03

R topics documented:

cfs.lps	2
contiguous.zoo	2
Cs	3
dC.dt	4
DOTemp	5
fmt.chron	5
Kt	6
lps.cfs	7
lps.cms	7
ODobbins	8
read.production	9
simp	10
SM	11
sunrise.set	13
window_chron	14

Index**16**

<code>cfs.lps</code>	<i>Convert from cubic feet per second to liters per second</i>
----------------------	--

Description

Convert from cubic feet persecond to liters per second

Usage

`cfs.lps(x)`

Arguments

`x` Discharge in cfs

Author(s)

Stephen A Sefick Jr.

<code>contiguous.zoo</code>	<i>contiguous.zoo</i>
-----------------------------	-----------------------

Description

find continuous non NA portions of zoo time series data

Usage

`contiguous.zoo(x)`

Arguments

`x` zoo time series object whatever indexes you want

Details

if you want to just find the contiguous portions of just one signal and not the interaction between two just duplicate the signal `contiguous.zoo(data.frame(x, coredata(x)))` should give you what you want

Examples

```
#single temperature
temp <- sample(20:30, 1)
Cs(temp)

#USGS Data (D0Temp)
library(chron)
library(zoo)
data(D0Temp)
Cs(D0Temp[, 1])
```

dC.dt

Change in Oxygen per time step

Description

Calculate the rate of change of Dissolved Oxygen

Usage

```
dC.dt(x)
```

Arguments

x Dissolved Oxygen time series

Details

input zoo series takes the difference of $DO_{t+1} - DO_t$

Value

zoo series of Dissolved Oxygen Differences with an NA for the first value as there is no value before that to subtract

Author(s)

Stephen A Sefick Jr.

References

Odum, H. T. (1956). "Primary production in flowing waters." *Limnology and Oceanography*, 1: 102-117.

Examples

```
data(D0Temp)
Diffconc <- dC.dt(D0Temp[, 2])
plot(Diffconc)
```

DOTemp	<i>Rhode River Street Pier–Maryland data set</i>
--------	--

Description

Test data set from Stephens and Jennings SWProd calculator (USGS). The data has been interpolated to make it have readings every fifteen minutes.

Usage

```
data(DOTemp)
```

Format

DateTime DateTime

Temp Temperature in Celcius

DO Dissolved Oxygen

Details

This is only for example and should be used judiciously for any kind of ecosystem interpretation (requires zoo and chron packages).

Source

Stephens, D.W., and Jennings, M.E., 1976, Determination of primary productivity and community metabolism in streams and lakes using diel oxygen measurements: U.S. Geological Survey Computer Contribution, 100 p.

fmt.chron	<i>Format Dates</i>
-----------	---------------------

Description

Used in the FUN argument of read.zoo for dates in the format mm/dd/yyyy hh:mm:ss

Usage

```
fmt.chron(x)
```

Arguments

x Data Time Column

Details

used internally in read.production

Author(s)

Stephen A Sefick Jr

See Also

[read.production](#)

Kt

Temperature Correction For Reaeration Value

Description

Temperature Correction For Reaeration Value. Corrects reaeration value to temperature of the stream.

Usage

Kt(K, temp)

Arguments

K	Rearation Coefficient single value or in a zoo object
temp	temperature value at time t in Degrees Celcius

Value

Single Values or zoo series

Author(s)

Stephen A Sefick Jr

References

Thyssen, N., Erlandsen, M., Jeppesen, E., Holm T. F., 1983. Modelling the reaeration capacity of low-land streams dominated by submerged macrophytes. In: Lauenroth, W.K., Skogerboe, G.V., Flug, M. (Eds.), *Analysis of Ecological Systems: State of the Art in Ecological Modelling*. Elsevier, pp. 861-867 as reported in Izagirre O., M. Bermejo, J. Pozo, and A. Elosegi. 2007. RIVERMET: An Excel-based tool to calculate river metabolism from diel oxygen-concentration curves. *Environmental Modelling & Software*, 22:24-32.

Examples

```
#data USGS
data(DOTemp)
#velocity 0.6, depth 0.4572
d <- ODobbins(0.6, 0.4572)
Kcorr <- Kt(d , DOTemp[,1])
```

lps.cfs

Liters Per Second to Cubic Feet per Second

Description

Convenience Function for converting from liters per second to cubic feet per second

Usage

```
lps.cfs(x)
```

Arguments

x Discharge in Liters per Second

Value

Discharge in Cubic feet per second

Author(s)

Stephen A Sefick Jr

Examples

```
lps.cfs(134000)
```

lps.cms

Liters per second to cubic meters per second

Description

Conversion Function

Usage

```
lps.cms(x)
```

Arguments

x discharge in Liters per second

Details

single value or if zoo series - zoo object

Value

single value or if zoo series - zoo object

Author(s)

Stephen A Sefick Jr.

Examples

```
lps.cms(134000)
```

ODobbins

*O'Conner Dobbins Surface Renewal Method for calculating Rearra-
tion Coeffiecient*

Description

calculate reaeration coefficient with the O'Conner Dobbins method

Usage

```
ODobbins(vel, dep)
```

Arguments

vel velocity in m/s
dep depth in meters

Details

Surface Renewal

Value

Reaeration Coefficient (1/d)

Author(s)

Stephen A Sefick Jr.

References

O'Connor, D. J., and W. E. Dobbins (1958). Mechanisms of reaeration in natural streams. Transactions of American Society of Civil Engineers, 123: 641-666.

Examples

```
#velocity 0.6
#depth 0.4572
ODobbins(0.6, 0.4572)
```

read.production	<i>Read in Time Series Data as zoo Object</i>
-----------------	---

Description

Wrapper Function to read.zoo

Usage

```
read.production(data)
```

Arguments

data	a csv file with headers and the date as mm/dd/yyyy hh:mm:ss format (think excel spreadsheet date format and the file is saved as a csv file)
------	--

Details

This is a wrapper function to read.zoo with a specific format required see above

Value

zoo object

Author(s)

Stephen A Sefick Jr

See Also

[read.table](#)

`simp`*Numeric Integration Using Simpson's method*

Description

Numeric Integration using the Simpson Method

Usage

```
simp(y, a = NULL, b = NULL, x = NULL, n = 200)
```

Arguments

<code>y</code>	y values to integrate
<code>x</code>	x values to integrate over
<code>a</code>	NULL
<code>b</code>	NULL
<code>n</code>	number of divisions defaults to 200

Value

Numeric Value of the integration

Author(s)

Rolf Turner

Examples

```
# 4-x^2-y^2
fun <- function(x, y){
  a <- 4
  b <- x^2
  d <- y^2
  z <- a-b-d
  return(z)
}

a <- fun(seq(-1000,1000,1), seq(-1000,1000,1))
simp(a, x=-1000:1000, n=1000)
```

Description

Calculate ER, NEP, and GPP from diurnal oxygen curves.

Usage

```
SM(depth, min_interval, DO, temp,
K, day, sr="00:00:00", ss="23:45:00",
start="00:00:00", end="23:45:00")
```

Arguments

depth	depth m K
min_interval	time resolution (e.g., 15 min)
temp	Zoo time Series temperature in degrees Celcius (see details)
DO	Zoo time Series Dissolved Oxygen in mg/L (see details)
day	date of the day of interest must be in quotes
start	time of the start of the "day" usually 00:00:00 must be in quotes
end	time of the end of the "day" usually 23:45:00 must be in quotes
sr	time of sunrise in the form 04:22:00 must be in quotes
ss	time of sunset in the form 19:23:00 must be in quotes
K	K at 20 deg. C (1/dt; e.g., 1/15min)

Details

The input data has to be a zoo time series constructed with a chron date time object of month/day/year hr:min:sec (i.e., 08/18/70 23:15:00)

sr and ss should be after and before the start and end of the time series, respectively.

ER is calculated as $\sum Et$ (i.e., mean nighttime NEP corrected for the difference in daytime temp and average nighttime temp)

GPP is calculated by summing NEP-ERT from sunrise to sunset

$NEP=ER+GPP$

Tested Against Rivermet spreadsheet (Izagirre 2007). The data from station 1 (7/10 - 7/15/2003) were used with $K=0.07$ from "Introduced K". ER, NEP, and GPP are in $mg/L*d$. The results were not identical. When Estimation from rivermet was regressed on estimation from this software, GPP, ER, and NEP intercepts did not differ significantly from 0 and slopes were nearly 1: 0.94, 0.91, and 0.95, respectively. Further testing is greatly appreciated.

Value

ER	Ecosystem Respiration
NEP	Net Ecosystem Production
GPP	Gross Primary Productivity

Author(s)

Stephen A Sefick Jr.

References

Odum, H. T. (1956). "Primary production in flowing waters." *Limnology and Oceanography*, 1: 102-117.

Thyssen, N., Erlandsen, M., Jeppesen, E., Holm T. F., 1983. Modelling the reaeration capacity of low-land streams

M.R. Grace and S.J. Imberger. 2006. "Stream Metabolism: Performing & Interpreting Measurements". Water Studies Centre Monash University, Murray Darling Basin Commission and New South Wales Department of Environment and Climate Change. 204 pp. Accessed at <http://www.sci.monash.edu.au/wsc/docs/manual-v3.pdf>

Izagirre, O., M. Bermejo, J. Pozo, and A. Elosegi. 2007. RIVERMET: An Excel-based tool to calculate river metabolism from diel oxygen concentration curves. *Environmental Modelling and Software*, 22: 24-32.

Examples

```
#zoo real data
#velocity 0.6
#depth 0.4572
#sunrise 6:00AM
#sunset 8:15PM
#K/96 to get K per dt (i.e., 96 15 min interval in 1 day)
data(DOTemp)

K <- ODobbins(0.6, 0.4572)
prod <- SM(min_interval=15, K=K/96,
depth=0.4572, temp=DOTemp[,1], DO=DOTemp[,2],
day="8/18/70", start="00:00:00",
end="23:45:00", sr="06:00:00", ss="20:15:00")
prod
```

`sunrise.set`*Calculate Sunrise Sunset Times*

Description

This function calculates sunrise sunset times in POSIXct and returns it in a handy dandy format to either export as a csv file or use directly in the calculation of Stream Metabolism. This function is based on maptools which is based on the NOAA sunrise sunset calculator.

Usage

```
sunrise.set(lat, long, date, timezone = "UTC", num.days = 1)
```

Arguments

<code>lat</code>	Latitude in decimal degrees
<code>long</code>	Longitude in decimal degrees
<code>date</code>	starting date (needs to be in quotes and in the format yyyy/mm/dd)
<code>timezone</code>	Time zone set to UTC default (needs to be in quotes)
<code>num.days</code>	1 if you just want only the calculation performed on "date" (default)

Details

Remember that the Prime Meridian is 0 through Greenwich, England. So anything W is - and anything E is +. Also anything in the Northern hemisphere is + latitude and Southern Hemisphere is - latitude. Generally UTC+5 is Eastern Standard Time, UTC+6 is CST, UTC+7 MST, UTC+8 PST. Another way of specifying time zones is Country City see examples. Be aware of timezones and daylight and standard time when using this function!!!!!! This will help you avoid headaches caused because minor oversites = large error in your calculations

Value

`output` data frame with all dates sunrise and sunset times specified

Author(s)

Stephen A Sefick Jr.

References

old site: <http://www.esrl.noaa.gov/gmd/grad/solcalc/sunrise.html>

new site: <http://www.esrl.noaa.gov/gmd/grad/solcalc/>

Examples

```
#This is for Atlanta Georgia
#(Only so that you can compare it to the NOAA
#website that is given above)
sunrise.set(33.43, -84.22, "2008/01/01", timezone="UTC+5")

#Same As above but look at Time Zone Specification
sunrise.set(33.43, -84.22, "2008/01/01", timezone="America/New_York")
```

window_chron	<i>Time Windows of Diurnal Curves</i>
--------------	---------------------------------------

Description

Takes a time window of a larger series

Usage

```
window_chron(x, day1, hour1, day2, hour2, ...)
```

Arguments

x	data to be subsetted
day1	start day
hour1	start time
day2	end date
hour2	end time
...	other arguments

Value

subset by time

Author(s)

Stephen A Sefick Jr.

References

chron, window, window.zoo

See Also

[window](#)

Examples

```
#with real data
data(DOTemp)
d <- window_chron(DOTemp, "8/18/70", "06:00:00", "8/18/70", "20:15:00")
plot(d)
```

Index

* **chron**

fmt.chron, 5
read.production, 9
window_chron, 14

* **datasets**

DOTemp, 5

* **math**

cfs.lps, 2
contiguous.zoo, 2
Cs, 3
dC.dt, 4
Kt, 6
lps.cfs, 7
lps.cms, 7
ODobbins, 8
simp, 10
SM, 11
sunrise.set, 13

cfs.lps, 2
contiguous.zoo, 2
Cs, 3

dC.dt, 4
DOTemp, 5

fmt.chron, 5

Kt, 6

lps.cfs, 7
lps.cms, 7

ODobbins, 8

read.production, 6, 9
read.table, 9

simp, 10
SM, 11
sunrise.set, 13

window, 14
window_chron, 14