

Package ‘bmixture’

September 11, 2019

Version 1.4

Date 2019-09-11

Title Bayesian Estimation for Finite Mixture of Distributions

Author Reza Mohammadi <<https://orcid.org/0000-0001-9538-0648>>

Maintainer Reza Mohammadi <a.mohammadi@uva.nl>

URL <https://www.uva.nl/profile/a.mohammadi>

Description Provides statistical tools for Bayesian estimation for finite mixture of distributions, mainly mixture of Gamma, Normal and t-distributions. The package is implemented the recent improvements in Bayesian literature for the finite mixture of distributions, including Mohammadi and et al. (2013) <[doi:10.1007/s00180-012-0323-3](https://doi.org/10.1007/s00180-012-0323-3)> and Mohammadi and Salehi-Rad (2012) <[doi:10.1080/03610918.2011.588358](https://doi.org/10.1080/03610918.2011.588358)>.

Depends R (>= 3.0.0)

Imports BDgraph

License GPL (>= 2)

Repository CRAN

NeedsCompilation yes

Date/Publication 2019-09-11 08:40:02 UTC

R topics documented:

bmixture-package	2
bmixgamma	3
bmixnorm	6
bmixt	9
galaxy	11
mixgamma	12
mixnorm	14
mixt	15
plot.bmixgamma	17
plot.bmixnorm	18
plot.bmixt	19

print.bmixgamma	20
print.bmixnorm	21
print.bmixt	22
rdirichlet	23
summary.bmixgamma	23
summary.bmixnorm	24
summary.bmixt	25

Index	27
--------------	-----------

bmixture-package	<i>Bayesian Estimation for Finite Mixture of Distributions</i>
------------------	--

Description

The R package **bmixture** provides statistical tools for Bayesian estimation in finite mixture of distributions. The package implemented the improvements in the Bayesian literature, including Mohammadi and Salehi-Rad (2012) and Mohammadi et al. (2013). Besides, the package contains several functions for simulation and visualization, as well as a real dataset taken from the literature.

How to cite this package

Whenever using this package, please cite as

Mohammadi R. (2019). **bmixture**: Bayesian Estimation for Finite Mixture of Distributions, R package version 1.3, <https://CRAN.R-project.org/package=bmixture>

Author(s)

Reza Mohammadi <a.mohammadi@uva.nl>

References

- Mohammadi, A., Salehi-Rad, M. R., and Wit, E. C. (2013) Using mixture of Gamma distributions for Bayesian analysis in an M/G/1 queue with optional second service. *Computational Statistics*, 28(2):683-700
- Mohammadi, A, and Salehi-Rad, M. R. (2012) Bayesian inference and prediction in an M/G/1 with optional second service. *Communications in Statistics-Simulation and Computation*, 41(3):419-435
- Stephens, M. (2000) Bayesian analysis of mixture models with an unknown number of components-an alternative to reversible jump methods. *Annals of statistics*, 28(1):40-74
- Richardson, S. and Green, P. J. (1997) On Bayesian analysis of mixtures with an unknown number of components. *Journal of the Royal Statistical Society: series B*, 59(4):731-792
- Green, P. J. (1995) Reversible jump Markov chain Monte Carlo computation and Bayesian model determination. *Biometrika*, 82(4):711-732
- Cappe, O., Christian P. R., and Tobias, R. (2003) Reversible jump, birth and death and more general continuous time Markov chain Monte Carlo samplers. *Journal of the Royal Statistical Society: Series B*, 65(3):679-700
- Wade, S. and Ghahramani, Z. (2018) Bayesian Cluster Analysis: Point Estimation and Credible Balls (with Discussion). *Bayesian Analysis*, 13(2):559-626

Examples

```

## Not run:

library( bmixture )

data( galaxy )

# Runing bdmcmc algorithm for the galaxy dataset
mcmc_sample = bmixnorm( data = galaxy )

summary( mcmc_sample )
plot( mcmc_sample )
print( mcmc_sample)

# simulating data from mixture of Normal with 3 components
n      = 500
mean   = c( 0 , 10 , 3 )
sd     = c( 1 , 1 , 1 )
weight = c( 0.3, 0.5, 0.2 )

data = rmixnorm( n = n, weight = weight, mean = mean, sd = sd )

# plot for simulation data
hist( data, prob = TRUE, nclass = 30, col = "gray" )

x      = seq( -20, 20, 0.05 )
densmixnorm = dmixnorm( x, weight, mean, sd )

lines( x, densmixnorm, lwd = 2 )

# Runing bdmcmc algorithm for the above simulation data set
bmixnorm.obj = bmixnorm( data, k = 3, iter = 1000 )

summary( bmixnorm.obj )

## End(Not run)

```

bmixgamma

Sampling algorithm for mixture of distributions

Description

This function consists of several sampling algorithms for Bayesian estimation for finite mixture of Gamma distributions.

Usage

```

bmixgamma( data, k = "unknown", iter = 1000, burnin = iter / 2, lambda = 1,
           mu = NULL, nu = NULL, kesi = NULL, tau = NULL, k.start = NULL,
           alpha.start = NULL, beta.start = NULL, pi.start = NULL,
           k_max = 30, trace = TRUE )

```

Arguments

data	The vector of data with size n.
k	The number of components of mixture distribution. Default is "unknown". It can take an integer values.
iter	The number of iteration for the sampling algorithm.
burnin	The number of burn-in iteration for the sampling algorithm.
lambda	For the case k = "unknown", it is the parameter of the prior distribution of number of components k.
mu	The parameter of alpha in mixture distribution.
nu	The parameter of alpha in mixture distribution.
kesi	The parameter of beta in mixture distribution.
tau	The parameter of beta in mixture distribution.
k.start	For the case k = "unknown", initial value for number of components of mixture distribution.
alpha.start	Initial value for parameter of mixture distribution.
beta.start	Initial value for parameter of mixture distribution.
pi.start	Initial value for parameter of mixture distribution.
k_max	For the case k = "unknown", maximum value for the number of components of mixture distribution.
trace	Logical: if TRUE (default), tracing information is printed.

Details

Sampling from finite mixture of Gamma distribution, with density:

$$Pr(x|k, \underline{\pi}, \underline{\alpha}, \underline{\beta}) = \sum_{i=1}^k \pi_i \text{Gamma}(x|\alpha_i, \beta_i),$$

where k is the number of components of mixture distribution (as a default we assume is unknown) and

$$\text{Gamma}(x|\alpha_i, \beta_i) = \frac{(\beta_i)^{\alpha_i}}{\Gamma(\alpha_i)} x^{\alpha_i-1} e^{-\beta_i x}.$$

The prior distributions are defined as below

$$P(K = k) \propto \frac{\lambda^k}{k!}, \quad k = 1, \dots, k_{max},$$

$$\pi_i|k \sim \text{Dirichlet}(1, \dots, 1),$$

$$\alpha_i|k \sim \text{Gamma}(\nu, \nu),$$

$$\beta_i|k \sim G(\eta, \tau),$$

for more details see Mohammadi et al. (2013).

Value

An object with S3 class "bmixgamma" is returned:

all_k	A vector which includes the waiting times for all iterations. It is needed for monitoring the convergence of the BD-MCMC algorithm.
all_weights	A vector which includes the waiting times for all iterations. It is needed for monitoring the convergence of the BD-MCMC algorithm.
pi_sample	A vector which includes the MCMC samples after burn-in from parameter pi of mixture distribution.
alpha_sample	A vector which includes the MCMC samples after burn-in from parameter alpha of mixture distribution.
beta_sample	A vector which includes the MCMC samples after burn-in from parameter beta of mixture distribution.
data	original data.

Author(s)

Reza Mohammadi <a.mohammadi@uva.nl>

References

- Mohammadi, A., Salehi-Rad, M. R., and Wit, E. C. (2013) Using mixture of Gamma distributions for Bayesian analysis in an M/G/1 queue with optional second service. *Computational Statistics*, 28(2):683-700
- Mohammadi, A. and Salehi-Rad, M. R. (2012) Bayesian inference and prediction in an M/G/1 with optional second service. *Communications in Statistics-Simulation and Computation*, 41(3):419-435
- Stephens, M. (2000) Bayesian analysis of mixture models with an unknown number of components—an alternative to reversible jump methods. *Annals of statistics*, 28(1):40-74
- Richardson, S. and Green, P. J. (1997) On Bayesian analysis of mixtures with an unknown number of components. *Journal of the Royal Statistical Society: series B*, 59(4):731-792
- Green, P. J. (1995) Reversible jump Markov chain Monte Carlo computation and Bayesian model determination. *Biometrika*, 82(4):711-732
- Cappe, O., Christian P. R., and Tobias, R. (2003) Reversible jump, birth and death and more general continuous time Markov chain Monte Carlo samplers. *Journal of the Royal Statistical Society: Series B*, 65(3):679-700
- Wade, S. and Ghahramani, Z. (2018) Bayesian Cluster Analysis: Point Estimation and Credible Balls (with Discussion). *Bayesian Analysis*, 13(2):559-626

See Also

[bmixnorm](#), [bmixt](#), [bmixgamma](#)

Examples

```
## Not run:
# simulating data from mixture of gamma with two components
n      = 1000 # number of observations
weight = c( 0.6, 0.4 )
alpha  = c( 12 , 1   )
beta   = c( 3  , 2   )

data <- rmixgamma( n = n, weight = weight, alpha = alpha, beta = beta )

# plot for simulation data
hist( data, prob = TRUE, nclass = 50, col = "gray" )

x      = seq( 0, 10, 0.05 )
truth  = dmixgamma( x, weight, alpha, beta )

lines( x, truth, lwd = 2 )

# Running bdmcmc algorithm for the above simulation data set
bmixgamma.obj <- bmixgamma( data, iter = 1000 )

summary( bmixgamma.obj )

plot( bmixgamma.obj )

## End(Not run)
```

 bmixnorm

Sampling algorithm for mixture of distributions

Description

This function consists of several sampling algorithms for Bayesian estimation for finite mixture of Normal distributions.

Usage

```
bmixnorm( data, k = "unknown", iter = 1000, burnin = iter / 2, lambda = 1,
          k.start = NULL, mu.start = NULL, sig.start = NULL, pi.start = NULL,
          k_max = 30, trace = TRUE )
```

Arguments

data	The vector of data with size n.
k	The number of components of mixture distribution. Default is "unknown". It can take an integer values.
iter	The number of iteration for the sampling algorithm.
burnin	The number of burn-in iteration for the sampling algorithm.

lambda	For the case k = "unknown", it is the parameter of the prior distribution of number of components k.
k.start	For the case k = "unknown", initial value for number of components of mixture distribution.
mu.start	Initial value for parameter of mixture distribution.
sig.start	Initial value for parameter of mixture distribution.
pi.start	Initial value for parameter of mixture distribution.
k_max	For the case k = "unknown", maximum value for the number of components of mixture distribution.
trace	Logical: if TRUE (default), tracing information is printed.

Details

Sampling from finite mixture of Gamma distribution, with density:

$$Pr(x|k, \underline{\pi}, \underline{\mu}, \underline{\sigma}) = \sum_{i=1}^k \pi_i N(x|\mu_i, \sigma_i),$$

where k is the number of components of mixture distribution (as a default we assume is unknown). The prior distributions are defined as below

$$P(K = k) \propto \frac{\lambda^k}{k!}, \quad k = 1, \dots, k_{max},$$

$$\pi_i|k \sim Dirichlet(1, \dots, 1),$$

$$\alpha_i|k \sim Gamma(\nu, \nu),$$

$$\beta_i|k \sim G(\eta, \tau),$$

for more details see Mohammadi et al. (2013) and Mohammadi and Salehi-Rad (2012).

Value

An object with S3 class "bmixnorm" is returned:

all_k	A vector which includes the waiting times for all iterations. It is needed for monitoring the convergence of the BD-MCMC algorithm.
all_weights	A vector which includes the waiting times for all iterations. It is needed for monitoring the convergence of the BD-MCMC algorithm.
pi_sample	A vector which includes the MCMC samples after burn-in from parameter pi of mixture distribution.
mu_sample	A vector which includes the MCMC samples after burn-in from parameter mu of mixture distribution.
sig_sample	A vector which includes the MCMC samples after burn-in from parameter sig of mixture distribution.
data	The original data.

Author(s)

Reza Mohammadi <a.mohammadi@uva.nl>

References

- Stephens, M. (2000) Bayesian analysis of mixture models with an unknown number of components-an alternative to reversible jump methods. *Annals of statistics*, 28(1):40-74
- Richardson, S. and Green, P. J. (1997) On Bayesian analysis of mixtures with an unknown number of components. *Journal of the Royal Statistical Society: series B*, 59(4):731-792
- Green, P. J. (1995) Reversible jump Markov chain Monte Carlo computation and Bayesian model determination. *Biometrika*, 82(4):711-732
- Cappe, O., Christian P. R., and Tobias, R. (2003) Reversible jump, birth and death and more general continuous time Markov chain Monte Carlo samplers. *Journal of the Royal Statistical Society: Series B*, 65(3):679-700
- Mohammadi, A., Salehi-Rad, M. R., and Wit, E. C. (2013) Using mixture of Gamma distributions for Bayesian analysis in an M/G/1 queue with optional second service. *Computational Statistics*, 28(2):683-700
- Mohammadi, A. and Salehi-Rad, M. R. (2012) Bayesian inference and prediction in an M/G/1 with optional second service. *Communications in Statistics-Simulation and Computation*, 41(3):419-435
- Wade, S. and Ghahramani, Z. (2018) Bayesian Cluster Analysis: Point Estimation and Credible Balls (with Discussion). *Bayesian Analysis*, 13(2):559-626

See Also

[bmixt](#), [bmixgamma](#), [rmixnorm](#)

Examples

```
## Not run:
data( galaxy )

# Running bdmcmc algorithm for the galaxy dataset
mcmc_sample = bmixnorm( data = galaxy )

summary( mcmc_sample )
plot( mcmc_sample )
print( mcmc_sample)

# simulating data from mixture of Normal with 3 components
n      = 500
weight = c( 0.3, 0.5, 0.2 )
mean   = c( 0 , 10 , 3 )
sd     = c( 1 , 1 , 1 )

data = rmixnorm( n = n, weight = weight, mean = mean, sd = sd )

# plot for simulation data
hist( data, prob = TRUE, nclass = 30, col = "gray" )
```



```

x          = seq( -20, 20, 0.05 )
densmixnorm = dmixnorm( x, weight, mean, sd )

lines( x, densmixnorm, lwd = 2 )

# Runing bdcmc algorithm for the above simulation data set
bmixnorm.obj = bmixnorm( data, k = 3, iter = 1000 )

summary( bmixnorm.obj )

## End(Not run)

```

 bmixt

Sampling algorithm for mixture of distributions

Description

This function consists of several sampling algorithms for Bayesian estimation for finite mixture of Normal distributions.

Usage

```

bmixt( data, k = "unknown", iter = 1000, burnin = iter / 2, lambda = 1, df = 1,
       k.start = NULL, mu.start = NULL, sig.start = NULL, pi.start = NULL,
       k_max = 30, trace = TRUE )

```

Arguments

data	The vector of data with size n.
k	The number of components of mixture distribution. Default is "unknown". It can take an integer values.
iter	The number of iteration for the sampling algorithm.
burnin	The number of burn-in iteration for the sampling algorithm.
lambda	For the case k = "unknown", it is the parameter of the prior distribution of number of components k.
df	Degrees of freedom (> 0, maybe non-integer). df = Inf is allowed.
k.start	For the case k = "unknown", initial value for number of components of mixture distribution.
mu.start	Initial value for parameter of mixture distribution.
sig.start	Initial value for parameter of mixture distribution.
pi.start	Initial value for parameter of mixture distribution.
k_max	For the case k = "unknown", maximum value for the number of components of mixture distribution.
trace	Logical: if TRUE (default), tracing information is printed.

Details

Sampling from finite mixture of Gamma distribution, with density:

$$Pr(x|k, \underline{\pi}, \underline{\mu}, \underline{\sigma}) = \sum_{i=1}^k \pi_i N(x|\mu_i, \sigma_i),$$

where k is the number of components of mixture distribution (as a default we assume is unknown). The prior distributions are defined as below

$$P(K = k) \propto \frac{\lambda^k}{k!}, \quad k = 1, \dots, k_{max},$$

$$\pi_i | k \sim \text{Dirichlet}(1, \dots, 1),$$

$$\alpha_i | k \sim \text{Gamma}(\nu, \nu),$$

$$\beta_i | k \sim G(\eta, \tau),$$

for more details see Mohammadi et al. (2013) and Mohammadi and Salehi-Rad (2012).

Value

An object with S3 class "bmixt" is returned:

all_k	A vector which includes the waiting times for all iterations. It is needed for monitoring the convergence of the BD-MCMC algorithm.
all_weights	A vector which includes the waiting times for all iterations. It is needed for monitoring the convergence of the BD-MCMC algorithm.
pi_sample	A vector which includes the MCMC samples after burn-in from parameter pi of mixture distribution.
mu_sample	A vector which includes the MCMC samples after burn-in from parameter mu of mixture distribution.
sig_sample	A vector which includes the MCMC samples after burn-in from parameter sig of mixture distribution.
data	The original data.

Author(s)

Reza Mohammadi <a.mohammadi@uva.nl>

References

- Stephens, M. (2000) Bayesian analysis of mixture models with an unknown number of components- an alternative to reversible jump methods. *Annals of statistics*, 28(1):40-74
- Richardson, S. and Green, P. J. (1997) On Bayesian analysis of mixtures with an unknown number of components. *Journal of the Royal Statistical Society: series B*, 59(4):731-792
- Green, P. J. (1995) Reversible jump Markov chain Monte Carlo computation and Bayesian model determination. *Biometrika*, 82(4):711-732

Cappe, O., Christian P. R., and Tobias, R. (2003) Reversible jump, birth and death and more general continuous time Markov chain Monte Carlo samplers. *Journal of the Royal Statistical Society: Series B*, 65(3):679-700

Mohammadi, A., Salehi-Rad, M. R., and Wit, E. C. (2013) Using mixture of Gamma distributions for Bayesian analysis in an M/G/1 queue with optional second service. *Computational Statistics*, 28(2):683-700

Mohammadi, A. and Salehi-Rad, M. R. (2012) Bayesian inference and prediction in an M/G/1 with optional second service. *Communications in Statistics-Simulation and Computation*, 41(3):419-435

Wade, S. and Ghahramani, Z. (2018) Bayesian Cluster Analysis: Point Estimation and Credible Balls (with Discussion). *Bayesian Analysis*, 13(2):559-626

See Also

[bmixnorm](#), [bmixgamma](#), [rmixt](#)

Examples

```
## Not run:
# simulating data from mixture of Normal with 3 components
n      = 500
weight = c( 0.3, 0.5, 0.2 )
mean   = c( 0 , 10 , 3 )
sd     = c( 1 , 1 , 1 )

data = rmixnorm( n = n, weight = weight, mean = mean, sd = sd )

# plot for simulation data
hist( data, prob = TRUE, nclass = 30, col = "gray" )

x      = seq( -20, 20, 0.05 )
densmixnorm = dmixnorm( x, weight, mean, sd )

lines( x, densmixnorm, lwd = 2 )

# Runing bdmcmc algorithm for the above simulation data set
bmixt.obj = bmixt( data, k = 3, iter = 1000 )

summary( bmixt.obj )

## End(Not run)
```

galaxy

Galaxy data

Description

This dataset considers of 82 observatons of the velocities (in 1000 km/second) of distant galaxies diverging from our own, from six well-separated conic sections of the Corona Borealis. The dataset has been analyzed under a variety of mixture models; See e.g. Stephens (2000).

Usage

```
data( galaxy )
```

Format

A data frame with 82 observations on the following variable.

speed a numeric vector giving the speed of galaxies (in 1000 km/second).

References

Stephens, M. (2000) Bayesian analysis of mixture models with an unknown number of components- an alternative to reversible jump methods. *Annals of statistics*, 28(1):40-74

Examples

```
data( galaxy )

hist( galaxy, prob = TRUE, xlim = c( 0, 40 ), ylim = c( 0, 0.3 ), nclass = 20,
      col = "gray", border = "white" )

lines( density( galaxy ), col = "black", lwd = 2 )
```

 mixgamma

Mixture of Gamma distribution

Description

Random generation and density function for the finite mixture of Gamma distribution.

Usage

```
rmixgamma( n = 10, weight = 1, alpha = 1, beta = 1 )
```

```
dmixgamma( x, weight = 1, alpha = 1, beta = 1 )
```

Arguments

n	The number of samples required.
x	The vector of quantiles.
weight	The vector of probability weights, with length equal to number of components (k). This is assumed to sum to 1; if not, it is normalized.
alpha	The vector of non-negative parameters of the Gamma distribution.
beta	The vector of non-negative parameters of the Gamma distribution.

Details

Sampling from finite mixture of Gamma distribution, with density:

$$Pr(x|\underline{w}, \underline{\alpha}, \underline{\beta}) = \sum_{i=1}^k w_i \text{Gamma}(x|\alpha_i, \beta_i),$$

where

$$\text{Gamma}(x|\alpha_i, \beta_i) = \frac{(\beta_i)^{\alpha_i}}{\Gamma(\alpha_i)} x^{\alpha_i-1} e^{-\beta_i x}.$$

Value

Generated data as an vector with size n .

Author(s)

Reza Mohammadi <a.mohammadi@uva.nl>

References

Mohammadi, A., Salehi-Rad, M. R., and Wit, E. C. (2013) Using mixture of Gamma distributions for Bayesian analysis in an M/G/1 queue with optional second service. *Computational Statistics*, 28(2):683-700

Mohammadi, A., and Salehi-Rad, M. R. (2012) Bayesian inference and prediction in an M/G/1 with optional second service. *Communications in Statistics-Simulation and Computation*, 41(3):419-435

See Also

[rmixnorm](#), [rmixt](#)

Examples

```
## Not run:
n      = 10000
weight = c( 0.6 , 0.3 , 0.1 )
alpha  = c( 100 , 200 , 300 )
beta   = c( 100/3, 200/4, 300/5 )

data = rmixgamma( n = n, weight = weight, alpha = alpha, beta = beta )

hist( data, prob = TRUE, nclass = 30, col = "gray" )

x      = seq( -20, 20, 0.05 )
densmixgamma = dmixnorm( x, weight, alpha, beta )

lines( x, densmixgamma, lwd = 2 )

## End(Not run)
```

 mixnorm

Mixture of Normal distribution

Description

Random generation and density function for the finite mixture of univariate Normal distribution.

Usage

```
rmixnorm( n = 10, weight = 1, mean = 0, sd = 1 )
```

```
dmixnorm( x, weight = 1, mean = 0, sd = 1 )
```

Arguments

n	The number of samples required.
x	The vector of quantiles.
weight	The vector of probability weights, with length equal to number of components (k). This is assumed to sum to 1; if not, it is normalized.
mean	The vector of means.
sd	The vector of standard deviations.

Details

Sampling from finite mixture of Normal distribution, with density:

$$Pr(x|\underline{w}, \underline{\mu}, \underline{\sigma}) = \sum_{i=1}^k w_i N(x|\mu_i, \sigma_i).$$

Value

Generated data as an vector with size n .

Author(s)

Reza Mohammadi <a.mohammadi@uva.nl>

References

- Mohammadi, A., Salehi-Rad, M. R., and Wit, E. C. (2013) Using mixture of Gamma distributions for Bayesian analysis in an M/G/1 queue with optional second service. *Computational Statistics*, 28(2):683-700
- Mohammadi, A., and Salehi-Rad, M. R. (2012) Bayesian inference and prediction in an M/G/1 with optional second service. *Communications in Statistics-Simulation and Computation*, 41(3):419-435

See Also

[rmixt](#), [rmixgamma](#)

Examples

```
## Not run:
n      = 10000
weight = c( 0.3, 0.5, 0.2 )
mean   = c( 0 , 10 , 3 )
sd     = c( 1 , 1 , 1 )

data = rmixnorm( n = n, weight = weight, mean = mean, sd = sd )

hist( data, prob = TRUE, nclass = 30, col = "gray" )

x      = seq( -20, 20, 0.05 )
densmixnorm = dmixnorm( x, weight, mean, sd )

lines( x, densmixnorm, lwd = 2 )

## End(Not run)
```

mixture

Mixture of t-distribution

Description

Random generation and density function for the finite mixture of univariate t-distribution.

Usage

```
rmixt( n = 10, weight = 1, df = 1, mean = 0, sd = 1 )
dmixt( x, weight = 1, df = 1, mean = 0, sd = 1 )
```

Arguments

n	The number of samples required.
x	The vector of quantiles.
weight	The vector of probability weights, with length equal to number of components (k). This is assumed to sum to 1; if not, it is normalized.
df	The vector of degrees of freedom (> 0 , maybe non-integer). $df = \text{Inf}$ is allowed.
mean	The vector of means.
sd	The vector of standard deviations.

Details

Sampling from finite mixture of t-distribution, with density:

$$Pr(x|\underline{w}, \underline{\mu}, \underline{\sigma}) = \sum_{i=1}^k w_i N(x|\mu_i, \sigma_i).$$

Value

Generated data as an vector with size n .

Author(s)

Reza Mohammadi <a.mohammadi@uva.nl>

References

Mohammadi, A., Salehi-Rad, M. R., and Wit, E. C. (2013) Using mixture of Gamma distributions for Bayesian analysis in an M/G/1 queue with optional second service. *Computational Statistics*, 28(2):683-700

Mohammadi, A., and Salehi-Rad, M. R. (2012) Bayesian inference and prediction in an M/G/1 with optional second service. *Communications in Statistics-Simulation and Computation*, 41(3):419-435

See Also

[rmixnorm](#), [rmixgamma](#)

Examples

```
## Not run:
n      = 10000
weight = c( 0.3, 0.5, 0.2 )
df     = c( 4 , 4 , 4 )
mean   = c( 0 , 10 , 3 )
sd     = c( 1 , 1 , 1 )

data = rmixt( n = n, weight = weight, df = df, mean = mean, sd = sd )

hist( data, prob = TRUE, nclass = 30, col = "gray" )

x      = seq( -20, 20, 0.05 )
densmixt = dmixt( x, weight, df, mean, sd )

lines( x, densmixt, lwd = 2 )

## End(Not run)
```

plot.bmixgamma *Plot function for S3 class "bmixgamma"*

Description

Visualizes the results for function [bmixgamma](#).

Usage

```
## S3 method for class 'bmixgamma'  
plot( x, ... )
```

Arguments

x An object of S3 class "bmixgamma", from function [bmixgamma](#).
... System reserved (no specific usage).

Author(s)

Reza Mohammadi <a.mohammadi@uva.nl>

See Also

[bmixgamma](#)

Examples

```
## Not run:  
# simulating data from mixture of gamma with two components  
n        = 500 # number of observations  
weight = c( 0.6, 0.4 )  
alpha  = c( 12 , 1    )  
beta   = c( 3  , 2    )  
  
data <- rmixgamma( n = n, weight = weight, alpha = alpha, beta = beta )  
  
# plot for simulation data  
hist( data, prob = TRUE, nclass = 50, col = "gray" )  
  
x        = seq( 0, 10, 0.05 )  
truth = dmixgamma( x, weight, alpha, beta )  
  
lines( x, truth, lwd = 2 )  
  
# Runing bdmcmc algorithm for the above simulation data set  
bmixgamma.obj <- bmixgamma( data )  
  
plot( bmixgamma.obj )  
  
## End(Not run)
```

plot.bmixnorm *Plot function for S3 class "bmixnorm"*

Description

Visualizes the results for function [bmixnorm](#).

Usage

```
## S3 method for class 'bmixnorm'  
plot( x, ... )
```

Arguments

x An object of S3 class "bmixnorm", from function [bmixnorm](#).
... System reserved (no specific usage).

Author(s)

Reza Mohammadi <a.mohammadi@uva.nl>

See Also

[bmixnorm](#)

Examples

```
## Not run:  
# simulating data from mixture of Normal with 3 components  
n        = 500  
weight = c( 0.3, 0.5, 0.2 )  
mean    = c( 0 , 10 , 3 )  
sd      = c( 1 , 1 , 1 )  
  
data = rmixnorm( n = n, weight = weight, mean = mean, sd = sd )  
  
# plot for simulation data  
hist( data, prob = TRUE, nclass = 30, col = "gray" )  
  
x                    = seq( -20, 20, 0.05 )  
densmixnorm = dmixnorm( x, weight, mean, sd )  
  
lines( x, densmixnorm, lwd = 2 )  
  
# Running bdmcmc algorithm for the above simulation data set  
bmixnorm.obj = bmixnorm( data, k = 3 )  
  
plot( bmixnorm.obj )  
  
## End(Not run)
```

`plot.bmixt`*Plot function for S3 class "bmixt"*

Description

Visualizes the results for function `bmixt`.

Usage

```
## S3 method for class 'bmixt'  
plot( x, ... )
```

Arguments

`x` An object of S3 class "bmixt", from function `bmixt`.
`...` System reserved (no specific usage).

Author(s)

Reza Mohammadi <a.mohammadi@uva.nl>

See Also

`bmixt`

Examples

```
## Not run:  
# simulating data from mixture of Normal with 3 components  
n      = 500  
weight = c( 0.3, 0.5, 0.2 )  
mean   = c( 0 , 10 , 3 )  
sd     = c( 1 , 1 , 1 )  
  
data = rmixnorm( n = n, weight = weight, mean = mean, sd = sd )  
  
# plot for simulation data  
hist( data, prob = TRUE, nclass = 30, col = "gray" )  
  
x      = seq( -20, 20, 0.05 )  
densmixnorm = dmixnorm( x, weight, mean, sd )  
  
lines( x, densmixnorm, lwd = 2 )  
  
# Runing bdmcmc algorithm for the above simulation data set  
bmixt.obj = bmixt( data, k = 3 )  
  
plot( bmixt.obj )  
  
## End(Not run)
```

print.bmixgamma *Print function for S3 class "bmixgamma"*

Description

Prints the information about the output of function [bmixgamma](#).

Usage

```
## S3 method for class 'bmixgamma'  
print( x, ... )
```

Arguments

x An object of S3 class "bmixgamma", from function [bmixgamma](#).
... System reserved (no specific usage).

Author(s)

Reza Mohammadi <a.mohammadi@uva.nl>

See Also

[bmixgamma](#)

Examples

```
## Not run:  
# simulating data from mixture of gamma with two components  
n        = 500 # number of observations  
weight = c( 0.6, 0.4 )  
alpha  = c( 12 , 1    )  
beta   = c( 3   , 2   )  
  
data <- rmixgamma( n = n, weight = weight, alpha = alpha, beta = beta )  
  
# plot for simulation data  
hist( data, prob = TRUE, nclass = 50, col = "gray" )  
  
x        = seq( 0, 10, 0.05 )  
truth = dmixgamma( x, weight, alpha, beta )  
  
lines( x, truth, lwd = 2 )  
  
# Runing bdmcmc algorithm for the above simulation data set  
bmixgamma.obj <- bmixgamma( data, iter = 500 )  
  
print( bmixgamma.obj )  
  
## End(Not run)
```

print.bmixnorm	<i>Print function for S3 class "bmixnorm"</i>
----------------	---

Description

Prints the information about the output of function [bmixnorm](#).

Usage

```
## S3 method for class 'bmixnorm'  
print( x, ... )
```

Arguments

x	An object of S3 class "bmixnorm", from function bmixnorm .
...	System reserved (no specific usage).

Author(s)

Reza Mohammadi <a.mohammadi@uva.nl>

See Also

[bmixnorm](#)

Examples

```
## Not run:  
# simulating data from mixture of Normal with 3 components  
n      = 500  
weight = c( 0.3, 0.5, 0.2 )  
mean   = c( 0 , 10 , 3 )  
sd     = c( 1 , 1 , 1 )  
  
data = rmixnorm( n = n, weight = weight, mean = mean, sd = sd )  
  
# plot for simulation data  
hist( data, prob = TRUE, nclass = 30, col = "gray" )  
  
x      = seq( -20, 20, 0.05 )  
densmixnorm = dmixnorm( x, weight, mean, sd )  
  
lines( x, densmixnorm, lwd = 2 )  
  
# Runing bdmcmc algorithm for the above simulation data set  
bmixnorm.obj = bmixnorm( data, k = 3, iter = 1000 )  
  
print( bmixnorm.obj )  
  
## End(Not run)
```

print.bmixt *Print function for S3 class "bmixt"*

Description

Prints the information about the output of function [bmixt](#).

Usage

```
## S3 method for class 'bmixt'  
print( x, ... )
```

Arguments

x An object of S3 class "bmixt", from function [bmixt](#).
... System reserved (no specific usage).

Author(s)

Reza Mohammadi <a.mohammadi@uva.nl>

See Also

[bmixt](#)

Examples

```
## Not run:  
# simulating data from mixture of Normal with 3 components  
n        = 500  
weight = c( 0.3, 0.5, 0.2 )  
mean    = c( 0 , 10 , 3 )  
sd      = c( 1 , 1 , 1 )  
  
data = rmixnorm( n = n, weight = weight, mean = mean, sd = sd )  
  
# plot for simulation data  
hist( data, prob = TRUE, nclass = 30, col = "gray" )  
  
x                    = seq( -20, 20, 0.05 )  
densmixnorm = dmixnorm( x, weight, mean, sd )  
  
lines( x, densmixnorm, lwd = 2 )  
  
# Runing bdmcmc algorithm for the above simulation data set  
bmixt.obj = bmixt( data, k = 3, iter = 1000 )  
  
print( bmixt.obj )  
  
## End(Not run)
```

`rdirichlet`*Random generation for the Dirichlet distribution*

Description

Random generation from the Dirichlet distribution.

Usage

```
rdirichlet( n = 10, alpha = c( 1, 1 ) )
```

Arguments

<code>n</code>	The number of samples required.
<code>alpha</code>	The vector of shape parameters.

Details

The Dirichlet distribution is the multidimensional generalization of the beta distribution.

Value

A matrix with `n` rows, each containing a single Dirichlet random deviate.

Author(s)

Reza Mohammadi <a.mohammadi@uva.nl>

Examples

```
draws = rdirichlet( n = 500, alpha = c( 1, 1, 1 ) )
boxplot( draws )
```

`summary.bmixgamma`*Summary function for S3 class "bmixgamma"*

Description

Provides a summary of the results for function [bmixgamma](#).

Usage

```
## S3 method for class 'bmixgamma'
summary( object, ... )
```

Arguments

object An object of S3 class "bmixgamma", from function [bmixgamma](#).
... System reserved (no specific usage).

Author(s)

Reza Mohammadi <a.mohammadi@uva.nl>

See Also

[bmixgamma](#)

Examples

```
## Not run:  
# simulating data from mixture of gamma with two components  
n        = 500 # number of observations  
weight = c( 0.6, 0.4 )  
alpha  = c( 12 , 1    )  
beta   = c( 3   , 2   )  
  
data <- rmixgamma( n = n, weight = weight, alpha = alpha, beta = beta )  
  
# plot for simulation data  
hist( data, prob = TRUE, nclass = 50, col = "gray" )  
  
x       = seq( 0, 10, 0.05 )  
truth  = dmixgamma( x, weight, alpha, beta )  
  
lines( x, truth, lwd = 2 )  
  
# Runing bdmcmc algorithm for the above simulation data set  
bmixgamma.obj <- bmixgamma( data, iter = 500 )  
  
summary( bmixgamma.obj )  
  
## End(Not run)
```

summary.bmixnorm

Summary function for S3 class "bmixnorm"

Description

Provides a summary of the results for function [bmixnorm](#).

Usage

```
## S3 method for class 'bmixnorm'  
summary( object, ... )
```


Arguments

object An object of S3 class "bmixnorm", from function [bmixnorm](#).
 ... System reserved (no specific usage).

Author(s)

Reza Mohammadi <a.mohammadi@uva.nl>

See Also

[bmixnorm](#)

Examples

```
## Not run:
# simulating data from mixture of Normal with 3 components
n      = 500
weight = c( 0.3, 0.5, 0.2 )
mean   = c( 0 , 10 , 3 )
sd     = c( 1 , 1 , 1 )

data = rmixnorm( n = n, weight = weight, mean = mean, sd = sd )

# plot for simulation data
hist( data, prob = TRUE, nclass = 30, col = "gray" )

x      = seq( -20, 20, 0.05 )
densmixnorm = dmixnorm( x, weight, mean, sd )

lines( x, densmixnorm, lwd = 2 )

# Runing bdmcmc algorithm for the above simulation data set
bmixnorm.obj = bmixnorm( data, k = 3, iter = 1000 )

summary( bmixnorm.obj )

## End(Not run)
```

summary.bmixt

Summary function for S3 class "bmixt"

Description

Provides a summary of the results for function [bmixt](#).

Usage

```
## S3 method for class 'bmixt'
summary( object, ... )
```

Arguments

object An object of S3 class "bmixt", from function [bmixt](#).
... System reserved (no specific usage).

Author(s)

Reza Mohammadi <a.mohammadi@uva.nl>

See Also

[bmixt](#)

Examples

```
## Not run:  
# simulating data from mixture of Normal with 3 components  
n        = 500  
weight = c( 0.3, 0.5, 0.2 )  
mean    = c( 0 , 10 , 3 )  
sd      = c( 1 , 1 , 1 )  
  
data = rmixnorm( n = n, weight = weight, mean = mean, sd = sd )  
  
# plot for simulation data  
hist( data, prob = TRUE, nclass = 30, col = "gray" )  
  
x            = seq( -20, 20, 0.05 )  
densmixnorm = dmixnorm( x, weight, mean, sd )  
  
lines( x, densmixnorm, lwd = 2 )  
  
# Runing bdmcmc algorithm for the above simulation data set  
bmixt.obj = bmixt( data, k = 3, iter = 1000 )  
  
summary( bmixt.obj )  
  
## End(Not run)
```

Index

- *Topic **datasets**
 - galaxy, 11
 - *Topic **distribution**
 - mixgamma, 12
 - mixnorm, 14
 - mixt, 15
 - rdirichlet, 23
 - *Topic **hplot**
 - plot.bmixgamma, 17
 - plot.bmixnorm, 18
 - plot.bmixt, 19
 - *Topic **iteration**
 - bmixgamma, 3
 - bmixnorm, 6
 - bmixt, 9
 - *Topic **package**
 - bmixture-package, 2
 - *Topic **print**
 - print.bmixgamma, 20
 - print.bmixnorm, 21
 - print.bmixt, 22
 - *Topic **sampling algorithms**
 - bmixgamma, 3
 - bmixnorm, 6
 - bmixt, 9
- bmixnorm_fixed_k (bmixture-package), 2
- bmixnorm_unknown_k
(bmixture-package), 2
- bmixgamma, 3, 5, 8, 11, 17, 20, 23, 24
- bmixnorm, 5, 6, 11, 18, 21, 24, 25
- bmixt, 5, 8, 9, 19, 22, 25, 26
- bmixture-package, 2
- dmixgamma (mixgamma), 12
- dmixnorm (mixnorm), 14
- dmixt (mixt), 15
- galaxy, 11
- mixgamma, 12
- mixnorm, 14
- mixt, 15
- plot.bmixgamma, 17
- plot.bmixnorm, 18
- plot.bmixt, 19
- print.bmixgamma, 20
- print.bmixnorm, 21
- print.bmixt, 22
- rdirichlet, 23
- rmixgamma, 15, 16
- rmixgamma (mixgamma), 12
- rmixnorm, 8, 13, 16
- rmixnorm (mixnorm), 14
- rmixt, 11, 13, 15
- rmixt (mixt), 15
- summary.bmixgamma, 23
- summary.bmixnorm, 24
- summary.bmixt, 25