

Package ‘plpoisson’

June 4, 2020

Type Package

Title Prediction Limits for Poisson Distribution

Version 0.1.1

Date 2020-06-01

Author Valbona Bejleri [aut] (<<https://orcid.org/0000-0001-9828-968X>>),
Luca Sartore [aut, cre] (<<https://orcid.org/0000-0002-0446-1328>>),
Balgobin Nandram [aut] (<<https://orcid.org/0000-0002-3204-0301>>)

Maintainer Luca Sartore <drwolf85@gmail.com>

Description Prediction limits for the Poisson distribution
are produced from both frequentist and Bayesian viewpoints. Limiting results
are provided in a Bayesian setting with uniform, Jeffreys and gamma as prior
distributions. More details on the methodology are discussed in Bejleri and
Nandram (2018) <[doi:10.1080/03610926.2017.1373814](https://doi.org/10.1080/03610926.2017.1373814)>.

License GPL-3

NeedsCompilation yes

Repository CRAN

Date/Publication 2020-06-04 12:20:06 UTC

R topics documented:

plpoisson-package	2
poisBayes	3
poisJEFF	5
poiss	6
poisUNIF	8

Index	10
--------------	-----------

Description

Prediction limits for Poisson distribution are useful when quantifying the uncertainty associated with predicting the occurrences of real life phenomena. The **plpoisson** package provides a set of functions to compute prediction limits of the inferred Poisson distribution under both, frequentist and Bayesian frameworks.

For frequentist prediction a common approach is to estimate the parameter based on the observed data firstly, then, to predict based on the estimated parameter. Different from the common approach of frequentist prediction, this approach does not require the estimation of the parameter. In a Bayesian setting, the uniform, Jeffreys and gamma distributions are used as priors when deriving the predictive posterior distribution.

Details

Package: plpoisson
 Type: Package
 Version: 0.1.1
 Date: 2020-06-01
 License: GPL-3

For a complete list of exported functions, use `library(help = "plpoisson")`.

Author(s)

Valbona Bejleri, Luca Sartore and Balgobin Nandram

Maintainer: Luca Sartore <drwolf85@gmail.com>

References

Bejleri, V., & Nandram, B. (2018). Bayesian and frequentist prediction limits for the Poisson distribution. *Communications in Statistics-Theory and Methods*, **47**(17), 4254-4271.

Bejleri, V. (2005). *Bayesian Prediction Intervals for the Poisson Model, Noninformative Priors*, Ph.D. Dissertation, American University, Washington, DC.

Examples

```
## Loading the package
library(plpoisson)

## Setting quantities of interest
xobs <- rpois(1, 50) # Number of the observed occurrences
n <- 1               # Total number of the time windows of
```

```

# of size 's' observed in the past
s <- rgamma(1, 4, .567) # Fixed size of observed time windows
t <- rgamma(1, 3, .33) # Future time window
a <- 5 # Shape hyperparameter of a gamma prior
b <- 1.558 # Rate hyperparameter of a gamma prior

## Frequentist prediction limits
poiss(xobs, n, s, t)

## Bayesian prediction limits (with uniform prior)
poisUNIF(xobs, n, s, t)

## Bayesian prediction limits (with Jeffreys prior)
poisJEFF(xobs, n, s, t)

## Bayesian prediction limits (with gamma prior)
poisBayes(xobs, n, s, t, a, b)

```

poisBayes

Bayesian Prediction Limits for Poisson Distribution (Gamma Prior)

Description

The function provides the Bayesian prediction limits of a Poisson random variable derived based on a gamma prior. The resulting prediction bounds quantify the uncertainty associated with the predicted future number of occurrences in a time window of size t .

Usage

```
poisBayes(xobs, n, s, t, a, b, alpha = 0.05)
```

Arguments

xobs	a numeric value denoting the number of the observed occurrences.
n	a numeric value representing the total number of the time windows s in the past (observed time windows).
s	a numeric value corresponding to the fixed size (or average size) of the observed time windows.
t	a numeric value indicating the size of the future time window.
a	a positive real number denoting the shape hyperparameter of a gamma prior distribution.
b	a positive real number representing the rate hyperparameter of a gamma prior distribution.
alpha	a numeric value associated to the credible probability. By default $\alpha = 0.05$, thus an prediction interval at 95% will be returned.

Details

When the argument $b = \text{Inf}$, one can obtain prediction limits with uniform prior by setting the argument $a = 1$. Similarly, one can get the limits with a Jeffreys prior by setting the argument $a = 0$.

Prediction bounds are obtained through the binary search algorithm.

Value

A list containing the following components:

lower	An integer value representing the lower bound of the prediction limit.
upper	An integer value representing the upper bound of the prediction limit.

Author(s)

Valbona Bejleri, Luca Sartore and Balgobin Nandram

References

- Bejleri, V., & Nandram, B. (2018). Bayesian and frequentist prediction limits for the Poisson distribution. *Communications in Statistics-Theory and Methods*, **47**(17), 4254-4271.
- Bejleri, V. (2005). *Bayesian Prediction Intervals for the Poisson Model, Noninformative Priors*, Ph.D. Dissertation, American University, Washington, DC.
- Davis, C. H. (1969). The binary search algorithm. *American Documentation (pre-1986)*, **20**(2), 167.

See Also

[poiss](#), [poisJEFF](#), [poisUNIF](#)

Examples

```
# Loading the package
library(plpoisson)
set.seed(2020L)

# Number of observed time windows
n <- 555L

# Simulating a dataset
data <- cbind.data.frame(
  occ_obs = rpois(n, rgamma(n, 5.5, .5)),
  win_siz = rgamma(n, 1.44, .777)
)

## Bayesian prediction limits
## (with gamma prior)
poisBayes(sum(data$occ_obs), # Past occurrences
  nrow(data), # Total past time windows
  mean(data$win_siz), # Window size
```

```
333, # Size of future window
2, 2.22) # Hyper-parameters for gamma prior
```

poisJEFF

Bayesian Prediction Limits for Poisson Distribution (Jeffreys Prior)

Description

The function provides the Bayesian prediction limits of a Poisson random variable derived based on a Jeffreys prior. The resulting prediction bounds quantify the uncertainty associated to the predicted future number of occurrences in a time windows of size t .

Usage

```
poisJEFF(xobs, n, s, t, alpha = 0.05)
```

Arguments

<code>xobs</code>	a numeric value denoting the number of the observed occurrences.
<code>n</code>	a numeric value representing the total number of the time windows s in the past (observed time windows).
<code>s</code>	a numeric value corresponding to the fixed size (or average size) of the observed time windows.
<code>t</code>	a numeric value indicating the size of the future time window.
<code>alpha</code>	a numeric value associated to the credible probability. By default $\alpha = 0.05$, thus an prediction interval at 95% will be returned.

Details

The resulting limits are equivalent to those provided when running the function `poisBayes()` with arguments $a = 0$ and $b = \text{Inf}$.

Prediction bounds are obtained through the binary search algorithm.

Value

A list containing the following components:

<code>lower</code>	An integer value representing the lower bound of the prediction limit.
<code>upper</code>	An integer value representing the upper bound of the prediction limit.

Author(s)

Valbona Bejleri, Luca Sartore and Balgobin Nandram

References

- Bejleri, V., & Nandram, B. (2018). Bayesian and frequentist prediction limits for the Poisson distribution. *Communications in Statistics-Theory and Methods*, **47**(17), 4254-4271.
- Bejleri, V. (2005). *Bayesian Prediction Intervals for the Poisson Model, Noninformative Priors*, Ph.D. Dissertation, American University, Washington, DC.
- Davis, C. H. (1969). The binary search algorithm. *American Documentation (pre-1986)*, **20**(2), 167.

See Also

[poiss](#), [poisBayes](#), [poisUNIF](#)

Examples

```
# Loading the package
library(plpoisson)
set.seed(2020L)

# Number of observed time windows
n <- 555L

# Simulating a dataset
data <- cbind.data.frame(
  occ_obs = rpois(n, rgamma(n, 5.5, .5)),
  win_siz = rgamma(n, 1.44, .777)
)

## Bayesian prediction limits
## (with Jeffreys prior)
poisJEFF(sum(data$occ_obs), # Past occurrences
  nrow(data), # Total past time windows
  mean(data$win_siz), # Window size
  444) # Size of future window
```

poiss

Frequentist Prediction Limits for Poisson Distribution

Description

The function provides the frequentist prediction limits of a Poisson random variable. The resulting prediction bounds quantify the uncertainty associated to the predicted future number of occurrences in a time windows of size t .

Usage

```
poiss(xobs, n, s, t, alpha = 0.05)
```

Arguments

xobs	a numeric value denoting the number of the observed occurrences.
n	a numeric value representing the total number of the time windows s in the past (observed time windows).
s	a numeric value corresponding to the fixed size (or average size) of the observed time windows.
t	a numeric value indicating the size of the future time window.
alpha	a numeric value associated to the probability of prediction. By default alpha = 0.05, thus a prediction interval at 95% will be returned.

Details

Prediction bounds are obtained through the binary search algorithm.

Value

A list containing the following components:

lower	An integer value representing the lower bound of the prediction limit.
upper	An integer value representing the upper bound of the prediction limit.

Author(s)

Valbona Bejleri, Luca Sartore and Balgobin Nandram

References

- Bejleri, V., & Nandram, B. (2018). Bayesian and frequentist prediction limits for the Poisson distribution. *Communications in Statistics-Theory and Methods*, **47**(17), 4254-4271.
- Bejleri, V. (2005). *Bayesian Prediction Intervals for the Poisson Model, Noninformative Priors*, Ph.D. Dissertation, American University, Washington, DC.
- Davis, C. H. (1969). The binary search algorithm. *American Documentation (pre-1986)*, **20**(2), 167.

See Also

[poisBayes](#), [poisJEFF](#), [poisUNIF](#)

Examples

```
# Loading the package
library(plpoisson)
set.seed(2020L)

# Number of observed time windows
n <- 555L

# Simulating a dataset
```

```
data <- cbind.data.frame(
  occ_obs = rpois(n, rgamma(n, 5.5, .5)),
  win_siz = rgamma(n, 1.44, .777)
)

## Frequentist prediction limits
poiss(sum(data$occ_obs), # Past occurrences
      nrow(data), # Total past time windows
      mean(data$win_siz), # Window size
      3) # Size of future window
```

poisUNIF

*Bayesian Prediction Limits for Poisson Distribution (Uniform Prior)***Description**

The function provides the Bayesian prediction limits of a Poisson random variable derived based on a uniform prior. The resulting prediction bounds quantify the uncertainty associated to the predicted future number of occurrences in a time windows of size t .

Usage

```
poisUNIF(xobs, n, s, t, alpha = 0.05)
```

Arguments

xobs	a numeric value denoting the number of the observed occurrences.
n	a numeric value representing the total number of the time windows s in the past (observed time windows).
s	a numeric value corresponding to the fixed size (or average size) of the observed time windows.
t	a numeric value indicating the size of the future time window.
alpha	a numeric value associated to the credible probability. By default $\alpha = 0.05$, thus an prediction interval at 95% will be returned.

Details

The resulting limits are equivalent to those provided when running the function `poisBayes()` with arguments $a = 1$ and $b = \text{Inf}$.

Prediction bounds are obtained through the binary search algorithm.

Value

A list containing the following components:

lower	An integer value representing the lower bound of the prediction limit.
upper	An integer value representing the upper bound of the prediction limit.

Author(s)

Valbona Bejleri, Luca Sartore and Balgobin Nandram

References

Bejleri, V., & Nandram, B. (2018). Bayesian and frequentist prediction limits for the Poisson distribution. *Communications in Statistics-Theory and Methods*, **47**(17), 4254-4271.

Bejleri, V. (2005). *Bayesian Prediction Intervals for the Poisson Model, Noninformative Priors*, Ph.D. Dissertation, American University, Washington, DC.

Davis, C. H. (1969). The binary search algorithm. *American Documentation (pre-1986)*, **20**(2), 167.

See Also

[poiss](#), [poisJEFF](#), [poisBayes](#)

Examples

```
# Loading the package
library(plpoisson)
set.seed(2020L)

# Number of observed time windows
n <- 555L

# Simulating a dataset
data <- cbind.data.frame(
  occ_obs = rpois(n, rgamma(n, 5.5, .5)),
  win_siz = rgamma(n, 1.44, .777)
)

## Bayesian prediction limits
## (with uniform prior)
poisUNIF(sum(data$occ_obs), # Past occurrences
  nrow(data), # Total past time windows
  mean(data$win_siz), # Window size
  444) # Size of future window
```

Index

- *Topic **bayesian**
 - poisBayes, [3](#)
 - poisJEFF, [5](#)
 - poisUNIF, [8](#)
- *Topic **distribution**
 - plpoisson-package, [2](#)
 - poisBayes, [3](#)
 - poisJEFF, [5](#)
 - poiss, [6](#)
 - poisUNIF, [8](#)
- *Topic **frequentist**
 - poiss, [6](#)
- *Topic **gamma prior**
 - poisBayes, [3](#)
- *Topic **intervals**
 - poisBayes, [3](#)
 - poisJEFF, [5](#)
 - poiss, [6](#)
 - poisUNIF, [8](#)
- *Topic **interval**
 - plpoisson-package, [2](#)
- *Topic **jeffreys prior**
 - poisJEFF, [5](#)
- *Topic **package**
 - plpoisson-package, [2](#)
- *Topic **poisson**
 - plpoisson-package, [2](#)
 - poisBayes, [3](#)
 - poisJEFF, [5](#)
 - poiss, [6](#)
 - poisUNIF, [8](#)
- *Topic **prediction**
 - plpoisson-package, [2](#)
 - poisBayes, [3](#)
 - poisJEFF, [5](#)
 - poiss, [6](#)
 - poisUNIF, [8](#)
- *Topic **uniform prior**
 - poisUNIF, [8](#)
- plpoisson (plpoisson-package), [2](#)
- plpoisson-package, [2](#)
- poisBayes, [3](#), [6](#), [7](#), [9](#)
- poisJEFF, [4](#), [5](#), [7](#), [9](#)
- poiss, [4](#), [6](#), [6](#), [9](#)
- poisUNIF, [4](#), [6](#), [7](#), [8](#)